



REFERENCE ONLY

## UNIVERSITY OF LONDON THESIS

Degree MD Year 2007 Name of Author DAWAS, Khaled

### COPYRIGHT

This is a thesis accepted for a Higher Degree of the University of London. It is an unpublished typescript and the copyright is held by the author. All persons consulting this thesis must read and abide by the Copyright Declaration below.

### COPYRIGHT DECLARATION

I recognise that the copyright of the above-described thesis rests with the author and that no quotation from it or information derived from it may be published without the prior written consent of the author.

### LOANS

Theses may not be lent to individuals, but the Senate House Library may lend a copy to approved libraries within the United Kingdom, for consultation solely on the premises of those libraries. Application should be made to: Inter-Library Loans, Senate House Library, Senate House, Malet Street, London WC1E 7HU.

### REPRODUCTION

University of London theses may not be reproduced without explicit written permission from the Senate House Library. Enquiries should be addressed to the Theses Section of the Library. Regulations concerning reproduction vary according to the date of acceptance of the thesis and are listed below as guidelines.

- A. Before 1962. Permission granted only upon the prior written consent of the author. (The Senate House Library will provide addresses where possible).
- B. 1962-1974. In many cases the author has agreed to permit copying upon completion of a Copyright Declaration.
- C. 1975-1988. Most theses may be copied upon completion of a Copyright Declaration.
- D. 1989 onwards. Most theses may be copied.

***This thesis comes within category D.***



This copy has been deposited in the Library of UCL



This copy has been deposited in the Senate House Library,  
Senate House, Malet Street, London WC1E 7HU.



# **Selective Endothelin Receptor Involvement in the Development of Colorectal Cancer and Liver Metastases**

**Khaled I Dawas MB BChir, MA, FRCS (Gen)**

A thesis submitted for the degree of Doctor of Medicine (MD)

University of London 2007

Department of Surgery

The Royal Free and University College Medical School, UCL, London

UMI Number: U592806

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U592806

Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author.  
Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against  
unauthorized copying under Title 17, United States Code.



ProQuest LLC  
789 East Eisenhower Parkway  
P.O. Box 1346  
Ann Arbor, MI 48106-1346

## STATEMENT OF ORIGINALITY

The studies described and presented in this thesis are the original work of the author. All work was performed by the author with the following exceptions:

- All autoradiography was performed in conjunction with Dr Michael Dashwood in Clinical Biochemistry, The Royal Free and University College Medical School, London.
- All flow cytometry was performed in conjunction with Dr Derek Davies at the Imperial Cancer Research Fund (Cancer Research UK).
- All specimens of colorectal liver metastases were collected, with patient consent, from surgical resections carried out by Professor I Taylor at The Middlesex Hospital, London.

No portion of the work referred to in this thesis has been submitted in support of an application for another degree or qualification of this or any other university, or other institute of learning.

All animal studies were performed under appropriate Home Office project and personal licences.

# CONTENTS

TITLE PAGE	1
STATEMENT OF ORIGINALITY	2
TABLE OF CONTENTS	3
ABSTRACT	6
ACKNOWLEDGMENTS	7
LIST OF ABBREVIATIONS	8
LIST OF FIGURES	10
LIST OF TABLES	11
CHAPTER I : REVIEW OF LITERATURE	12
1.1 ENDOTHELINS	13
1.1.1 INTRODUCTION	13
1.1.2 STRUCTURE	14
1.1.3 GENE REGULATION	14
1.1.4 LOCAL PRODUCTION	15
1.1.5 CLEARANCE FROM PLASMA	15
1.1.6 ENDOTHELIN-1	15
1.1.7 ENDOTHELIN-2 & ENDOTHELIN-3	16
1.2 ENDOTHELIN RECEPTORS	16
1.2.1 LOCALISATION & FUNCTION	17
1.2.2 ENDOTHELIN RECEPTOR NOMENCLATURE & SUBCLASSIFICATION	18
1.2.3 CELLULAR MECHANISMS	19
1.3 ENDOTHELIN ANTAGONISTS	20
1.4 ENDOTHELIN-1 & CARDIOVASCULAR DISEASE	23
1.5 ENDOTHELIN-1 & CANCER	24
1.5.1 THE ROLE OF ANGIOGENESIS IN CANCER DEVELOPMENT	25

1.5.2 ENDOTHELIN-1 & ITS ROLE IN TUMOUROGENESIS	28
1.6 COLORECTAL CANCER	30
1.7 LIVER METASTASES	32
1.7.1 SUMMARY	32
1.7.2 DETECTION & IMAGING OF LIVER METASTASES	34
1.7.3 CURRENT MANAGEMENT OF COLORECTAL LIVER METASTASES – SURGERY	35
1.7.4 OTHER THERAPIES	39
1.8 AIMS	42
CHAPTER II : MECHANISMS OF ENDOTHELIN-1 STIMULATED PROLIFERATION IN CANCER CELLS	43
2.1 INTRODUCTION	44
2.2 MATERIALS & METHODS	45
2.2.1 CELL CULTURE	45
2.2.2 ET-1 ASSAYS	45
2.2.3 GROWTH ASSAYS	46
2.2.4 IMMUNOHISTOCHEMISTRY	47
2.2.5 FLOW CYTOMETRY	48
2.2.6 STATISTICAL ANALYSIS	48
2.3 RESULTS	49
2.3.1 ET-1 ASSAYS	49
2.3.2 GROWTH ASSAYS	49
2.3.3 IMMUNOHISTOCHEMISTRY	56
2.3.4 FLOW CYTOMETRY	57
2.4 DISCUSSION	59

<b>CHAPTER III: ENDOTHELIN RECEPTOR EXPRESSION IN COLORECTAL LIVER METASTASES</b>	<b>66</b>
<b>3.1 INTRODUCTION</b>	<b>67</b>
<b>3.2 MATERIALS &amp; METHODS</b>	<b>68</b>
<b>3.2.1 TISSUE COLLECTION AND PREPARATION</b>	<b>68</b>
<b>3.2.2 IN VITRO AUTORADIOGRAPHY</b>	<b>68</b>
<b>3.2.3 DOUBLE LABELLING</b>	<b>69</b>
<b>3.2.4 STATISTICAL ANALYSIS</b>	<b>70</b>
<b>3.3 RESULTS</b>	<b>70</b>
<b>3.3.1 IN VITRO AUTORADIOGRAPHY</b>	<b>70</b>
<b>3.3.2 DOUBLE LABELLING</b>	<b>71</b>
<b>3.4 DISCUSSION</b>	<b>73</b>
 <b>CHAPTER IV: THE EFFECT OF AN ORAL ET<sub>A</sub> RECEPTOR ANTAGONIST ON COLORECTAL LIVER METASTASES IN AN ANIMAL MODEL</b>	 <b>78</b>
<b>4.1 INTRODUCTION</b>	<b>79</b>
<b>4.2 METHODS</b>	<b>80</b>
<b>4.2.1 CELL CULTURE</b>	<b>80</b>
<b>4.2.2 ANIMALS</b>	<b>80</b>
<b>4.2.3 INOCULATION OF CANCER CELLS</b>	<b>81</b>
<b>4.2.4 TISSUE SPECIMENS</b>	<b>81</b>
<b>4.3 RESULTS</b>	<b>81</b>
<b>4.4 DISCUSSION</b>	<b>82</b>
 <b>CHAPTER V: SUMMARY AND CONCLUSION OF THESIS</b>	 <b>86</b>
 <b>BIBLIOGRAPHY</b>	 <b>90</b>



## **Abstract**

### **Selective Endothelin Receptor Involvement in the Development of Colorectal Cancer and Liver Metastases**

Endothelin-1 (ET-1), a vasoactive peptide with mitogenic qualities, is over-expressed in the plasma of patients with colorectal cancer (CRC) and liver metastases. Its actions are mediated via two G-protein linked receptors, ET<sub>A</sub> and ET<sub>B</sub>. This study (i) examined the effect of ET-1 and its receptor antagonists as well as a G-protein blocker on CRC cell line growth; (ii) investigated tET receptor distribution in human CRC and liver metastases by autoradiography; (iii) recorded the effect on tumour growth of an oral ET<sub>A</sub> receptor antagonist to an in vivo model of colorectal cancer liver metastases. ET-1 stimulated significant growth in the cancer cells. This effect was reversed by the ET<sub>A</sub>, but not the ET<sub>B</sub>, receptor antagonist. Apoptosis was similar in controls and ET-1± antagonist treated cells. BrDU staining demonstrated an ET-1 dependent increase in mitosis, reversed by the ET<sub>A</sub> receptor antagonist. Blocking the G-protein subunits also reversed growth. In human cancer tissues ET<sub>A</sub> was over-expressed while ET<sub>B</sub> was under-expressed compared to controls. There was no significant difference in weight or number of liver metastases between control and experimental rats in vivo. In summary, ET-1 stimulates human CRC growth in vitro via ET<sub>A</sub> receptors by mechanisms that include stimulation of mitosis but not alteration of apoptosis. Thi signal is transduced via the G-protein subunits Go or Gi. Human CRC liver metastases tissue over-express ET<sub>A</sub> receptors compared to normal tissue. ET receptor antagonists may have a therapeutic role in primary and metastatic CRC.

## **Acknowledgments**

Professor I Taylor and Professor G Burnstock directed and supported me throughout this project. For that and their enduring patience I am indebted to them. Dr M Loizidou was, and remains, a role model of dedication, selfless advice, academic acumen and most importantly to me, of a true friend.

I am also indebted to Dr M Dashwood for his patience and unreserved support and advice. His time and expertise made significant segments of this thesis possible.

Mr S Fredericks, at St George Hospital Medical School, and Dr D Davies, at the ICRF, both provided valuable technical support and I am very grateful for to them.

Last, but not least, the thanks of a son, brother and husband to my family and Rouba Sbano.

## List of Abbreviations

5-FU	5-fluorouracil
AJCC	The American Joint Committee on Cancer
APC	adenomatous polyposis coli
BrDU	5-bromo-2'-deoxyuridine
CEA	Carcinoembryonic antigen
COX-1/COX-2	Cyclo-oxygenase-1 /-2
CRC	Colorectal cancer
CT	Computerised tomography
DAB	3,3'-diaminobenzidine tetrahydrochloride
DMEM	Dulbecco's Modified Eagle Medium
EDTA	Ethylenediaminetetraacetic acid
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
ET	Endothelin
ET-1/ET-2/ET-3	Endothelin-1 / -2 / -3
FAP	Familial adenomatous polyposis
FCS	Foetal calf serum
FDG-PET	F-18 fluorodeoxyglucose PET
FUDR	Fluorodeoxyuridine
HAI	Hepatic artery infusion
HNPCC	Hereditary non-polyposis colorectal cancer
HPI	Hepatic portal infusion
IOUS	Intraoperative ultrasound

IP3	Inositol Triphosphate
MAPK	Mitogen activated protein kinase
MRI	Magnetic resonance image
NO	Nitric oxide
PBS	Phosphate buffered saline
PHR	Percentage hepatic replacement
PI3K	Phosphoinositide 3-kinase
PKC	Protein kinase C
PLC	Phospholipase C
PT	Pertussis toxin
RFA	Radiofrequency ablation
TGF- $\beta$	Tumour growth factor $\beta$
TNF- $\alpha$	Tumour necrosis factor $\alpha$
UICC	International Union Against Cancer
US	Ultrasound
VEGF	Vascular endothelial growth factor

## LIST OF FIGURES

<b>2.1 EFFECT OF ENDOTHELIN-1 ON GROWTH OF MC28</b>	<b>51</b>
<b>2.2 EFFECT OF ENDOTHELIN-1 ON GROWTH OF SW620</b>	<b>52</b>
<b>2.3 EFFECT OF ENDOTHELIN-1 ON GROWTH OF LIM1215</b>	<b>52</b>
<b>2.4 EFFECT OF ET-1 / ET<sub>A</sub> RECEPTOR ANTAGONIST ON GROWTH OF MC28 CELL</b>	<b>53</b>
<b>2.5 EFFECT OF ET-1 / ET<sub>A</sub> RECEPTOR ANTAGONIST ON GROWTH OF SW620 CELLS</b>	<b>54</b>
<b>2.6 EFFECT OF ET-1 / ET<sub>A</sub> RECEPTOR ANTAGONIST ON GROWTH OF LIM1215 CELLS</b>	<b>54</b>
<b>2.7 EFFECT OF ET<sub>A</sub> RECEPTOR ANTAGONISM ON GROWTH OF SW620 CELLS</b>	<b>55</b>
<b>2.8 EFFECT OF G-PROTEIN INHIBITION ON ET-1 STIMULATED CELLS</b>	<b>56</b>
<b>2.9 FLOW CYTOGRAMS FOR MC28 CELLS</b>	<b>58</b>
<b>3.1 ENDOTHELIN RECEPTOR EXPRESSION IN SPECIMENS OF NORMAL LIVER AND LIVER METASTASES</b>	<b>72</b>
<b>3.2 AUTORADIOGRAPHY IMAGE: ET RECEPTOR BINDING IN SECTIONS OF HUMAN LIVER METASTASES</b>	<b>72</b>
<b>3.3 DOUBLE LABELLING IMAGE – LIVER METASTASES</b>	<b>73</b>

## LIST OF TABLES

1.1 LIVER METASTASES STAGING SYSTEM	34
1.2 IMAGING INVESTIGATIONS USED IN THE DIAGNOSIS OF COLORECTAL LIVER METASTASES	36
1.3 FACTORS INFLUENCING SURVIVAL FOLLOWING RESECTION OF LIVER METASTASES	39
2.1 THE EFFECT OF ET-1 AND ET <sub>B</sub> RECEPTOR ANTAGONIST ON CELL GROWTH IN CULTURE	63
2.2 BRDU STAINING RESULTS IN CELLS STIMULATED BY ET-1 ± ET <sub>A</sub> ANTAGONIST	64
2.3 PROPORTION OF APOPTOTIC CELLS FOLLOWING EXPOSURE TO ET-1 ± ET <sub>A</sub> ANATGONIST	65
3.1 ENDOTHELIN RECEPTOR EXPRESSION IN SPECIMENS OF NORMAL LIVER AND LIVER METASTASES	71
4.1 RESULTS OF ANIMAL STUDIES	82

## **CHAPTER I**

### **Review of Literature**

Colorectal cancer is one of the commonest cancers in the West. Despite improvements in surgical and oncological treatment, survival at 5 years remains at 50% and one of the main factors for this poor outcome is the high rate of liver metastases. There is an increasing understanding of the underlying genetic and cellular biology of this disease. Endothelin-1 (ET-1) has a role to play in the development of colorectal cancer and its metastases. It may therefore be possible to use ET receptor antagonists in the treatment of colorectal cancer metastases. The introduction reviews current knowledge of ET-1 with special emphasis on its role in cancer generally and colorectal cancer specifically.

## **1.1 Endothelins**

### **1.1.1 Introduction**

A family of peptides found to cause coronary vasoconstriction in 1985 (Hickey *et al*, 1985) was subsequently isolated and identified in 1988 (Yanagisawa *et al*, 1988), and named endothelins (ET). There are three recognised members of this family - endothelin-1 (ET-1), endothelin-2 (ET-2) and endothelin-3 (ET-3) - which are 21 amino acid peptides and act as modulators of vasomotor tone, cell proliferation and hormone production in a variety of tissues. However, more recently Kishi (1998) described a 31 amino acid peptide with potent vasocontractile activity which was inhibited by both endothelin receptors A and B (ET<sub>A</sub> and ET<sub>B</sub>) specific antagonists. Collectively, the ET family ligands, precursors and associated enzymes and the ET receptors are called the "ET axis" (Nelson JB *et al*, 1999).



### 1.1.2 Structure

Preproendothelin-1 is 212 amino acids long and is processed to the 38 amino acid prohormone big endothelin-1, which is secreted and circulates in the plasma. The glycoprotein enzyme endothelin-converting enzyme (ECE)(Yanagisawa *et al*, 1988) cleaves big endothelin-1 between positions 21 (tryptophan) and 22 (valine) to generate ET-1.

### 1.1.3 Gene Regulation

The genes for ET-1, ET-2 and ET-3 have been localised to human chromosomes 6, 1 and 20 respectively (Bloch *et al*, 1989; 1991; Arinami *et al*, 1991). In general, the expression pattern of the ET genes is analogous to the tissue localisation pattern of ET receptors (Haynes *et al*, 1993) and suggests that ET is most likely a local mediator, acting as an autocrine or paracrine factor.

It is generally known that little, if any, ET-1 is stored in endothelial cells under physiological conditions and therefore must be made constitutively and then released from the cells. Endothelial cell expression of preproET-1 mRNA is known to be altered by a number of factors, including haemodynamic shear stress (Morita *et al*, 1993), hypoxia (Kourembanas *et al*, 1991), transforming growth factor- $\beta$  (TGF- $\beta$ )(Kurihara *et al*, 1989), tumour necrosis factor- $\alpha$  (TNF- $\alpha$ )(Marsden PA *et al*, 1992) and thrombin (Yanagisawa *et al*, 1988). Endothelin production is inhibited by agents that stimulate intracellular cGMP levels, including endothelium derived relaxing factor (EDRF), nitrovasodilators, prostaglandins and heparin (Saijonmaa *et al*, 1990; Boulanger *et al*, 1991; Kohno *et al*, 1992; 1993; Imai *et al*, 1993; Yokokawa *et al*, 1993)

#### **1.1.4 Local Production**

Endothelial cells appear to be the primary source of ET circulating in plasma. It is, however, possible that a variety of other cell types may release ET as happens in a wide range of cultured cells. The isoform of ET most prevalent in culture media and in body fluids or tissue is ET-1. ET-3 may be predominant in some tissues, particularly gut and brain. ET-2 has been difficult to detect by readily available methods due to cross-reactivity of most antibodies with ET-1 and therefore, little is known about it.

Localisation studies using molecular probes have shown that ET-1 is made where it is likely to have an effect, supporting the hypothesis that ET-1 is a paracrine factor. ET-1 produced by human umbilical vein endothelial cells and grown to confluent monolayers on acellular amniotic membranes has been shown to release predominantly into the basolateral compartment. ET-1 is thus released toward what would be the underlying vascular smooth muscle cells of the vessel intima (Wagner *et al*, 1992).

#### **1.1.5 Clearance from Plasma**

ET is thought to be removed from the circulation and extra-circulatory spaces by binding to its receptors (ET<sub>B</sub>) followed by internalisation, uncoupling and enzymatic degradation. Because of this high capacity clearance mechanism, free and intact ET would be expected to be scarce in biological tissue or fluid, as measurements of ET levels by immunoassay have demonstrated.

#### **1.1.6 Endothelin-1**

Endothelin-1 (ET-1) is produced in both endothelial and vascular smooth muscle

cells. As much as 75% of ET-1 secretion from cultured endothelial cells is towards the vascular smooth muscle side of the cells, where it can bind to specific receptors to cause vasoconstriction. While it is unlikely that this pool of ET-1 contributes to the plasma concentration it remains, nevertheless, useful to measure plasma endothelin-1 levels as they have been found to correlate well with severity of disease, such as in congestive cardiac failure (Wei C-M *et al*, 1994). ET-1 probably has a role in the maintenance of basal vasomotor tone. The infusion of BQ-123, a selective antagonist of ET<sub>A</sub> receptors, into the brachial artery of normal subjects caused progressive vasodilation and a 64% increase in forearm blood flow after one hour (Haynes WG *et al*, 1994). Although each of the endothelins is capable of producing vasoconstriction, ET-1 is the most potent vasoconstrictor known to man for it is 100 times more potent than noradrenaline on a molar basis. ET-1 also potentiates the vasoconstriction caused by noradrenaline (Tabuchi *et al*, 1989) and vice versa.

#### **1.1.7 Endothelin-2 and Endothelin-3**

Endothelin-2 (ET-2) is produced predominantly within the kidney and intestine but the cells of origin are unclear. It has no unique physiological functions as compared to ET-1.

Endothelin-3 (ET-3), like ET-1, circulates in the plasma. It has been found ubiquitously in the brain, the gastrointestinal tract, the kidneys and lungs.

### **1.2 Endothelin Receptors**

The endothelins bind to two types of receptors (A & B) in mammalian tissue. The

receptors are members of the super family of receptors linked with guanine-nucleotide-binding (G) proteins and range from 45,000 to 50,000 daltons in size in various tissues. The amino acid structure of the two types of receptor is approximately 50% identical.

The human ET<sub>A</sub> and ET<sub>B</sub> receptor genes have been assigned to chromosome 4 (Hosoda *et al*, 1992) and chromosome 13 (Arai *et al*, 1993) respectively. A third receptor type, ET<sub>C</sub>, which selectively responds to ET-3 has been identified and cloned in *Xenopus laevis* dermal melanophore (Karne *et al*, 1993). A mammalian equivalent has not been identified.

The greatest similarity between ET<sub>A</sub> and ET<sub>B</sub> receptors is in the trans-membrane regions and the intracellular loops. The greatest dissimilarities are found in the N-termini and extracellular loops 1 and 3. Alteration of the N-terminal extracellular region alters ET-1 binding to the human ET<sub>A</sub> receptor (Hashido *et al*, 1992) and results in differential binding affinity for the latter (ET-1>ET-2>ET-3).

### **1.2.1 Localisation and Function**

ET<sub>A</sub> is relatively abundant in human lung, aorta and heart and much less abundant in liver, pancreas, kidney, brain, adrenal, gut and skeletal muscle (Hosoda *et al*, 1992; Hori *et al*, 1992; Elshourbagy *et al*, 1993). ET<sub>B</sub> is relatively abundant in cerebral cortex, cerebellum, liver, kidney, lung and placenta and much less abundant in pancreas, gut, heart and skeletal muscle (Elshourbagy *et al*, 1993). In general, within the vascular wall, smooth muscle cells mainly express ET<sub>A</sub> and endothelial cells mainly express ET<sub>B</sub> receptors (Hosoda *et al*, 1991). However, this generalisation is somewhat controversial in the light of results from functional studies. Moreland (1994) concluded from their functional studies, using BQ-123 and

sarafotoxin S6c ( a selective ET<sub>B</sub> receptor agonist), that ET<sub>A</sub> receptors predominate on the high pressure (arterial) side of the circulation and ET<sub>B</sub> receptors predominate on the low pressure (venous) side of the circulation. Thus it seems clear that ET-1 can produce contraction in arteries (i.e., vascular smooth muscle cells) via either an ET<sub>A</sub> or ET<sub>B</sub>-mediated pathway.

### 1.2.2 Endothelin Receptor Nomenclature and Subclassification

Though only two mammalian endothelin receptors have been isolated, cloned and expressed to date, the existence of additional endothelin receptors and/or endothelin receptor subtypes have been postulated based on pharmacological studies that cannot be explained on the basis of only the ET<sub>A</sub> and ET<sub>B</sub> receptor.

**ET<sub>A</sub> receptor subtypes:** Initial work relating to the classification of endothelin responses suggested that endothelin-induced vasoconstriction was mediated by the ET<sub>A</sub> receptor situated on vascular smooth muscle cells. There are numerous examples demonstrating inhibition of ET-1 induced vasoconstriction and mitogenesis by the ET<sub>A</sub> receptor antagonist BQ-123 (Ihara M *et al*, 1992a). Karaki and colleagues (1993) observed that some ET<sub>A</sub> receptor mediated responses were insensitive to inhibition by BQ-123, and they therefore proposed the existence of ET<sub>A</sub> subtypes. It was postulated that the ET<sub>A1</sub> receptor represented a subtype which is sensitive to inhibition by BQ-123 while ET<sub>A2</sub> is insensitive to this type of inhibition.

**ET<sub>B</sub> receptor subtypes:** There are numerous examples of pharmacological data indicating the existence of anatomically separate and distinct ET<sub>B</sub> receptor subtypes; an ET<sub>B</sub> receptor on the endothelium that produces the release of nitric oxide to mediate vasodilation, and a subtype on vascular smooth muscle that directly mediates vasoconstriction (Warner TD *et al*, 1993a; b; Douglas SA *et al*, 1995). ET<sub>B1</sub>

was the name given to those  $ET_B$  receptors upon which the compound IRL1038 acted as selective antagonist and the compound PD142893 acted as a non-selective antagonist (Warner TD *et al*, 1993a). The  $ET_{B2}$  agonists were not affected by either of those compounds (Sudjarwo *et al*, 1993). The  $ET_{B1}$  receptors are in the endothelium and bring about the release of relaxing factors, while the  $ET_{B2}$  receptors are found in vascular smooth muscle and when activated cause vasoconstriction.

**Other endothelin receptors:** The  $ET_C$  receptor displays more affinity with ET-3 than with ET-1. It appears to trigger nitrous oxide (NO) release (Karne *et al*, 1993). Until now no homologous receptor has been discovered for it in mammal cells.

### 1.2.3 Cellular Mechanisms

Attempts have been made to explain the mechanism of action of ET on vascular smooth muscle. The vasoconstrictor response is affected by two types of receptor,  $ET_A$  and  $ET_{B2}$ . This vasoconstriction is correlated with a maintained heightening of intracellular calcium ( $Ca^{2+}$ ) produced via a two-phase mechanism. The binding of ET-1 to the receptor activates phospholipase C which hydrolyses phosphatidylinositol-4,5-bisphosphate (PIP<sub>2</sub>) into two products, inositol triphosphate (IP<sub>3</sub>) and diacylglycerol (DAG). The initial increase in  $Ca^{2+}$  concentration is caused by rapid mobilisation of intracellular  $Ca^{2+}$  stores by IP<sub>3</sub>. This initial amount of  $Ca^{2+}$  entering would be able to depolarise the vascular smooth muscle sufficiently for this in turn to activate the L-type voltage-dependent  $Ca^{2+}$  channels. Contraction could also be partially affected by the action of another product of PIP<sub>2</sub> hydrolysis, namely DAG.

In addition to the vasoactive effects, ET-1 promotes proliferation of vascular endothelial cells (via  $ET_B$ ) and of vascular smooth muscle (via  $ET_A$ ) and fibroblasts

(via both receptors) resulting in structural changes to the vessel wall. The mitogenic effect of ET-1 is attributed to stimulation of tyrosine residues as well as phosphorylation and activation of two other mitogen-activated protein kinases in the vascular smooth muscle (Golfman *et al*, 1993; Hirata *et al*, 1989; Bobik *et al*, 1990).

### 1.3 Endothelin Antagonists

Experimental and clinical evidence exists to support the importance of the development of drugs that block the production or actions of ET for use in cardiovascular medicine and oncology, particularly in conditions in which these peptides are clearly implicated.

The first ET receptor antagonists were of a peptide nature. In the development of these compounds, it was discovered that ET<sub>A</sub> receptors recognise the tertiary structure of the N-terminal and C-terminal portion of the ET molecules. However, ET<sub>B</sub> receptors only recognise the structure of the C-terminal portion. In 1991 based on these conclusions the first described ET antagonist, IRL1038, was developed; this was a selective in vivo inhibitor of ET<sub>B</sub> receptors capable of antagonising ET-1 and ET-3-induced contractions in guinea pig ileum (Sakurai *et al*, 1993). The first attempt at obtaining a selective ET<sub>A</sub> receptor antagonist was made by taking a fragment of ET peptide with a modification of the disulphide bridges and with the introduction of a diaminopropionic acid in the structure. The compound thereby obtained inhibited the ET-1-induced contraction, but not that induced by ET-3 in the pulmonary arteries (Spinella *et al*, 1991). Based on these studies the compound BQ-123 was introduced. The latter is now accepted to be the gold standard compound for ET receptor identification (Ihara *et al*, 1992). The potency of ET antagonists

varies according to what we consider to be their short or long term action. In this way it was discovered that BQ-123 totally blocks the production of IP<sub>3</sub> provoked by ET-1 in vascular smooth muscle cells, but not the mitogenic effect on them.

Competitiveness studies show that this loss of long term activity is not related to the degradation of the compound, but rather with dissociation processes. Another ET<sub>A</sub> receptor antagonist of a peptide nature which has been developed is the compound BQ-610. This compound antagonises cerebral vasoconstriction produced by subarachnoid haemorrhage, which is thought to be mediated by endogenous ET-1 acting through ET<sub>A</sub> receptors (Zuccarello *et al*, 1994).

Non-peptide ET antagonists have also been developed. The first was the compound 50-235 that was a selective ET<sub>A</sub> receptor agonist with no effect on ET<sub>B</sub> (Mihara *et al*, 1993). Some of the later compounds developed did not have heightened selectivity for one receptor or the other and were therefore considered to be ET<sub>A</sub>/ET<sub>B</sub> receptor antagonists. The best known non-peptide antagonist is the compound Ro-47-0203, also known as Bosentan. Bosentan shows 20 to 50-fold selectivity for the ET<sub>A</sub> receptor in studies of the inhibition of [<sup>125</sup>I]ET-1 binding to recombinant human receptors. In functional studies, however, it was somewhat less selective (Clozel *et al*, 1994). Other studies have shown that the systemic administration of Bosentan to patients with severe chronic heart failure produces systemic, pulmonary and peripheral venous vasodilation and improved cardiac performance, without causing reflex tachycardia (Kiowski *et al*, 1995). On the other hand it should be pointed out that the compound BQ-788 is a non-peptide antagonist selective for ET<sub>B</sub> receptors. This compound has been widely used in animal experiments for the characterisation of receptors which mediate the effects of ET. It can potentiate the arterial blood pressure increases induced by ET-1 and this is because it blocks the ET<sub>B</sub> receptors



in the endothelium and vascular smooth muscle; as a result, their initial vasodilator effect is sometimes lost (Ishikawa *et al*, 1994).

A novel class of orally active non-peptide ET<sub>A</sub> antagonists has been discovered by high throughput screening. The compounds are ET<sub>A</sub> selective, are orally available and show a long duration of action (Riechers *et al*, 1996). One of these is the compound A-127722, the most potent ET receptor antagonist disclosed to date, which has a more than 1000-fold greater affinity for binding to human ET<sub>A</sub> receptors than to human ET<sub>B</sub> receptors. It has been shown to be orally bio-available in the rat, dog and monkey. In the conscious rat, A-127722 exhibits a dose-related inhibition of the blood pressure response to exogenous ET-1 (Opgenorth *et al*, 1996).

The use of ET antagonists in cancer has demonstrated some benefit and further potential. Ali (2000b) demonstrated the mitogenic effect of ET-1 on human colorectal cancer cell lines *in vitro*. This action was inhibited by the application of the ET<sub>A</sub> antagonist BQ-123. Moraitis (1997) and a group led by Bagnato (1997) also demonstrated similar results whilst working on ovarian cancer cell lines *in vitro*. Similar results in prostate cancer cell lines underpin ongoing clinical trials using ET<sub>A</sub> antagonists in prostate cancer (Zonnenberg *et al*, 2003; Carducci *et al*, 2003). The proliferation of three human pancreatic carcinoma cell lines expressing ET<sub>A</sub> receptors was significantly reduced using a selective ET<sub>A</sub> antagonist (Bhargava S *et al*, 2005). *In vivo*, the selective ET<sub>A</sub> inhibition resulted in reduced angiogenesis as measured by lower microvessel densities (Bhargava *et al*, 2005). The biological mechanisms of ET-1 action in cancer are described further on in this thesis.

## 1.4 Endothelin-1 and Cardiovascular Disease

The endothelins are involved in the patho-physiological mechanism of a number of vascular conditions. Evidence now suggests a role for ET in sustained vasoconstriction as seen in hypertension and heart failure, local ischaemia as in myocardial infarction and acute renal failure and vasospasm as in Raynaud's disease and subarachnoid haemorrhage.

Possibly all isoforms of ET may be involved in the pathogenesis of hypertension through several effects. In addition to increasing central (Nishimura *et al*, 1991) and peripheral (Wong-Dusting *et al*, 1990) sympathetic activity, it stimulates the generation of renin, angiotensin II, aldosterone (Rakugi *et al*, 1990), as well as the generation of adrenaline (Boarder *et al*, 1989). Besides its vasoconstrictor and pressor effect, ET-1 has positive inotropic properties (Ishikawa *et al*, 1988; McClellan G *et al*, 1996) and an antinatriuretic action (Rabelink *et al*, 1994). In moderate to severe human essential hypertension the expression of ET-1 gene is increased though plasma levels of ET-1 are only slightly elevated in experimental and human hypertension. This may be due to the fact that ET synthesised in the endothelium is mostly released towards the underlying smooth muscle cells.

The plasma concentration of Big ET-1 correlates well with the severity of heart failure and is the most powerful predictor of outcome (Masson S *et al*, 2006).

Ischaemia can induce synthesis of endothelins. In humans, plasma endothelial levels are increased in acute myocardial infarction and unstable angina and the higher the plasma endothelin level in myocardial infarction and unstable angina the worse the prognosis (Miyauchi *et al*, 1989). Plasma ET levels on the third day after myocardial infarction are significantly related to mortality and plasma ET levels at 9

weeks after hospitalisation with unstable angina are significantly related to the incidence of further cardiovascular events (Omland T *et al*, 1994).

ET has been implicated in the acute renal failure and hypertension associated with nephrotoxic agents such as cyclosporin (Kon *et al*, 1990). The renal vasculature is very sensitive to ET and administration of ET antagonists 24 hrs after the ischaemic event still provided marked protection against severe acute renal failure (Gellai *et al*, 1995). ET-1 antagonists also play a role in the treatment of patients with pulmonary hypertension (Barst RJ *et al*, 2006).

ET is a potent vasoconstrictor of isolated cerebral arteries and its effects are potentiated after subarachnoid haemorrhage (Alafaci *et al*, 1990).

## **1.5 Endothelin-1 and Cancer**

ET-1 has been found to act as a mitogen for human epithelial carcinoma cell lines *in vitro* (Shichiri *et al*, 1991). The known ET-1 positive mitogenic effects affect fibroblasts and cancer cells (Kusuhara *et al*, 1989; 1990). Elevated levels of ET-1 have been found in specimens of cancer tissue including prostate (Nelson J *et al*, 1995), colon (Asham *et al*, 1997), breast (Patel *et al*, 1995) and thyroid (Donckier J *et al*, 2004). Patients with primary or metastatic colorectal and primary liver cancers are also known to have elevated plasma ET-1 levels (Shankar *et al*, 1996; Nakamuta *et al*, 1993). ET receptors are upregulated in specimens of prostate (Nelson J *et al*, 1995), lung (Zhao *et al*, 1995), breast (Wulfing P *et al*, 2004) and thyroid papillary (Donckier J *et al*, 2004) cancers. In *in vitro* systems, ovarian cancer cell lines produce ET-1 and possess ET<sub>A</sub> receptors (Bagnato *et al*, 1997). Bagnato, Moraitis and others, working on ovarian cancer cell lines, also found that addition of

exogenous ET resulted in specific dose-dependent increases in cell number for ET-1 and ET-2 but not for ET-3. In the presence and absence of the exogenous peptides, the addition of BQ-123 reduced the cell number. This result is consistent with ET-1 acting via ET<sub>A</sub> receptors as a positive mitogen (Moraitis *et al*, 1997; Bagnato *et al*, 1997) and is discussed further in section 1.5.2.

Tumourigenesis is dependent on the parallel development of a vascular supply (Folkman J, 1972). The vessels originate from the host circulation and their development correlates positively with the metastatic potential of the tumour (Folkman J, 1971; 1972). Recent studies have attempted to manipulate the vessel growth in several cancers with a view to inhibiting growth of the primary and subsequent metastases. ET-1, a recognised vasoconstrictor peptide, has been implicated as an angiogenic agent (Dawas *et al*, 1999, Knowles *et al*, 2005). With the recent development of ET-1 antagonists, the manipulation of angiogenesis within tumours may be possible.

### **1.5.1 The Role of Angiogenesis in Cancer Development**

Most solid human tumours will exist in situ for a substantial length of time without the need for vascularisation (Folkman J in *The Molecular Basis of Cancer*, 1995).

Vascularisation is necessary for progression of the neoplasm beyond 2-3mm<sup>3</sup> and its ability to metastasise (Folkman J, 1971; Weidner N, 1995). A “switch”, occurring in some of the tumour cells, to an angiogenic phenotype initiates vascularisation (Folkman J in *The Molecular Basis of Cancer*, 1995). This “switch” involves a change in the steady state equilibrium between positive and negative regulators of microvessel growth (Dameron *et al*, 1994).

The process of tumour angiogenesis starts with vessels and tumour growing towards each other (Harris A, 1997). Driven by angiogenic factors (bFGF, VEGF, tumour necrosis factor- $\beta$  and PDGF) and dampened by the down-regulation of certain angiogenic inhibitors (angiostatin, endostatin, platelet factor 4 and interleukin 1), these “driven” endothelial cells divide much more rapidly than normal endothelial cells. The endothelial basement membrane is degraded initiating new vessel sprouting. The new tumour vasculature is abnormal in that it is leaky, thus allowing passage of fibrinogen and plasminogen. Effectively, the normal process of angiogenesis is incomplete since maturation is halted. The regulatory imbalances may be the key to this derangement of the angiogenic process in tumours. The maturation would thus be inhibited at a primitive stage characterised by leaky vessel walls. Tsujii (1998) (working on *in vitro* model systems involving co-culture of endothelial cells with colon carcinoma cells) demonstrated that cyclo-oxygenase (a key enzyme required for the conversion of arachidonic acid to prostaglandins) plays an important role in the promotion of tumour related angiogenesis. The latter was inhibited by the application of non-steroidal anti-inflammatory compounds.

Neo-vascularisation will usually herald the onset of symptoms and tumours may then become clinically detectable yet it is at this stage that tumours become less accessible to chemotherapy (Folkman J, 1995). The tumours possess an increased interstitial pressure from the leaky tumour vessels and the absence of intra-tumour lymphatic drainage. Thus vascular compression and eventually central necrosis ensue. Chemotherapeutic agents are unable to penetrate the tumour, partly for this reason.

Growth and metastases of tumours are known to be dependent on neo-vascularisation for several reasons. A high degree of tumour vascularisation with

newly formed leaky vessels increases the chance for tumour cells to enter the circulatory system and metastasise (Folkman J, 1995). At the cellular level, neo-vascularisation augments tumour growth through a “perfusion” effect which allows metabolic exchange and a “paracrine” effect which supplies growth factors (Folkman J, 1971; Shalaby F *et al*, 1995). Perfusion is more efficient than diffusion in allowing metabolite and catabolite exchange in crowded tissue.

The angiogenic / tumourigenic effects of ET-1 could be mediated through the known mechanism of VEGF stimulation and enhanced endothelial mitogenesis in the existing tumour or vasculature. As mentioned earlier, selective ET<sub>A</sub> receptor inhibition *in vivo* resulted in reduced angiogenesis as measured by lower microvessel densities (Bhargava S *et al*, 2005). Wulfig (2004) observed a significant positive correlation between microvessel density and ET expression.

Moreover, expression of VEGF was found more often in tumours with over-expression of ET and its receptors. In ovarian cancer, ET-1 stimulated VEGF production through the ET<sub>A</sub> receptor and selective inhibition of this receptor significantly impaired the angiogenic response (Salani D *et al*, 2000)

Inhibition of VEGF has shown potent anti-tumour effects in *in vivo* animal models (Millauer *et al*, 1996). Thus ET-1 antagonists would play their anti tumour role as both anti-angiogenic and anti-mitogenic agents.

A number of anti-angiogenic therapies are proving effective in combating solid tumours. The focus of the therapies has comprised inhibiting positive angiogenesis regulators, enhancing negative ones or directly targeting tumour endothelial cells. The latter technique has proven effectiveness as demonstrated by Huang *et al* (1997) who demonstrated complete tumour regression of neuroblastoma xenografts in athymic mice. Similarly, Pipili-Synetos *et al*(1995), working on the *in vivo* model of

angiogenesis of the chick chorioallantoic membrane (CAM) and the model of growth and metastasis in the Lewis Lung carcinoma in mice, obtained results suggesting that angiogenesis inhibitors (isosorbide 5-mononitrate and isosorbide dinitrate) inhibited tumour growth and metastasis. Furthermore, in rodents, anti-angiogenic therapy increases the delivery of chemotherapy to a tumour (Teicher *et al*, 1994). This is explained by the suggestion that anti-angiogenic factors unpack the mass of tumour cells (Folkman, 1995), thus lowering the tumour's interstitial pressure. Recently, anti-VEGF agents have, indeed, improved prognosis in patients with metastatic colorectal cancer, when added to standard chemotherapy (Kabbinar FF *et al*, 2006).

### **1.5.2 Endothelin-1 and its Role in Tumourogenesis**

Raised levels of ET-1 in specimens of various cancers are well established. Patients with primary or metastatic colorectal and primary liver cancers are also known to have elevated plasma ET-1 levels (Shankar *et al*, 1996; Nakamuta *et al*, 1993). ET receptor density is also altered with ET<sub>A</sub> upregulated, as detected by autoradiography, in specimens of prostate, lung and colorectal cancers while ET<sub>B</sub> is downregulated (Ali *et al*, 2000a, Nelson J *et al*, 1995, Zhao *et al*, 1995).

There are several reports indicating correlation between ET-1 expression and prognosis in cancer. In breast cancer the expression of ET-1 and both its receptors correlated with acquisition of malignant potential (Wulfing P *et al*, 2003). Univariate analysis has shown that ET-1 expression, in addition to sex, nodal metastatic involvement and stage, were significant predictors of poor survival and disease free survival outcome figures in non-small cell lung carcinoma (Boldrini *et al*, 2005). In colorectal cancer, a Cox regression model identified age, advanced stage and

plasma big ET-1 levels as independent variables of significant prognostic value for overall survival (Arun *et al*, 2004).

Immunohistochemistry has shown that ET-1 is over-expressed in specimens of colorectal (Asham *et al*, 1997) hepatocellular (Kar *et al*, 1995) and prostate cancer (Nelson *et al*, 1996). A study conducted in our centre has also shown that colorectal cancer patients, with or without liver metastases, have elevated plasma levels of ET-1 (Shankar A *et al*, 1998). This is in agreement with studies on patients with hepatocellular cancer (Nakamuta *et al*, 1993). Also of interest is the finding that ET<sub>A</sub> receptors are upregulated in ovarian (Bagnato *et al*, 1999), colorectal (Ali H *et al*, 2000a), lung (Zhao *et al*, 1995) and meningioma / glioma (Harland *et al*, 1998) suggesting that autocrine or paracrine ET-1 loops contribute to cancer growth and progression (Shichiri *et al*, 1991; Moraitis *et al*, 1997). The frequent concomitant downregulation of ET<sub>B</sub> receptors in the same cancer types (Nelson J *et al*, 1996), suggests that ET<sub>B</sub> is not used for any cancer associated signals or even its apparent loss may remove some as yet unidentified inhibitory loop.

In vitro, several cancer cell lines have demonstrated proliferation in response to exogenous ET-1. The cell lines include colonic, ovarian, and meningioma (Kitagawa *et al*, 1994). This effect was reversed on further addition of an ET<sub>A</sub> receptor antagonist as is also the case in pancreatic cancer cell lines (Bhargava *et al*, 2005). Ovarian cancer cell lines produce ET-1 and possess ET<sub>A</sub> receptors (Bagnato *et al*, 1997). Moraitis (1997) working on two ovarian cancer cell lines, further found that addition of exogenous ET resulted in specific dose-dependent increases in cell number for ET-1 and ET-2 but not for ET-3. In the presence and absence of the exogenous peptides, the addition of BQ-123 reduced the cell number. This result is consistent with ET-1 acting via ET<sub>A</sub> receptors as a positive mitogen. Downstream



events which propagate the mitogenic effect on ovarian cancer cells include EGFR transactivation and MAPK (Bagnato *et al*, 1999). Overall the known ET-1 positive mitogenic effects are not restricted to endothelial cells but also affect fibroblasts and cancer cells (Kusuhara *et al*, 1989; 1990).

Specifically in CRC, my field of research, the application of an ET<sub>A</sub> antagonist reduced cell growth in at least one human colorectal cell line (Ali *et al*, 2000 b). A pilot study using a syngeneic rat model of liver metastases demonstrated that the intraportal administration of the ET<sub>A</sub> specific antagonist, BQ-123, shortly after tumour cells were inoculated via the same route, resulted in significant reduction in tumour load (Asham *et al*, 1997). These results may be due to the inhibition of the mitogenic effect of ET-1 but there is strong evidence for the anti-angiogenic effect complementing the former if not instigating it.

## **1.6 Colorectal Cancer**

Colorectal cancer is the second commonest cause of cancer death comprising 10% of all cancer deaths in the UK. In England and Wales there are 17,000 deaths each year attributed to colorectal cancer (mortality stats: reviews of the register general on deaths by cause, sex and age in England & Wales 1993 (revised 1994); Vukasin AP *et al*, 1990; Cancer Research UK 2005 <http://info.cancerresearchuk.org/cancerstats/types/bowel/incidence> ). Of the annual incidence of 30,000 new cases per year 10-12,000 are rectal cancers (Cancer Research Campaign. Factsheet 18.1 1993, 2005 Cancer Research UK <http://info.cancerresearchuk.org/cancerstats/types/bowel/incidence>). This distinction is important for deciding on neoadjuvant and adjuvant therapy and for the sake of comparisons between outcomes. Within the colon about 50% of cancers arise in the left side and 25% in the right (Cancer

Research UK 2005 [http://info.cancerresearchuk.org/cancerstats /types/bowel](http://info.cancerresearchuk.org/cancerstats/types/bowel)).

The aetiology of colorectal cancer is multifactorial. It is now accepted that there is an inherited and sporadic form. Approximately 5% of colorectal cancer cases diagnosed are attributable to an underlying hereditary colorectal cancer syndrome (Kwak EL *et al*, 2007). These include hereditary non-polyposis colorectal cancer (HNPCC) which is inherited in an autosomal dominant fashion. It arises due to mutations in mismatch repair genes, whose role is to repair mistakes in base pair matching during DNA replication. The clinical diagnosis of HNPCC families is based on the Amsterdam criteria I and II of the International Collaborative Group on HNPCC (ICG-HNPCC) (Vasen HFA *et al*, 1999). Familial adenomatous polyposis (FAP) is less common than HNPCC but is associated with a 100% risk of developing colorectal cancer. Again, it is inherited in an autosomal dominant fashion and is due to a mutation in the tumour suppressor adenomatous polyposis coli (APC) gene on chromosome 5q (Wu JS *et al*, 1998; Potter JD 1999). Other less common inherited colorectal cancers occur in patients with the autosomal dominant Peutz-Jeghers syndrome and juvenile polyposis.

Genetic aspects to the development of sporadic colorectal cancer tend to be similar to those found in FAP cancers and comprise mutations of the APC, K-ras and p53 genes. The latter mutations are found in more than 60% of all adenomas and carcinomas (Powell SM *et al*, 1992; Vogelstein B *et al*, 1988; Scott N *et al*, 1993; Kikuchi-Yanoshita *et al*, 1992). These mutations drive the adenoma carcinoma sequence. The remaining sporadic cancers are genetically closer to HNPCCs. There has been evidence that dietary factors such as lack of fibre and vegetables, and excess of animal fats are associated with an increased risk of colorectal cancer for some time now (Burkitt, 1971; Modan *et al*, 1975; Bjelke, 1973). There has also

been recent, reserved, enthusiasm about the inhibitory effect of COX-2 inhibitors on colonic adenoma and adenomatous polyp progression (Bertagnolli MM *et al*, 2006; Arber N *et al*, 2006).

Recent advances in colorectal cancer have focused on combinations of monoclonal therapies and chemotherapy which have resulted in better clinical outcomes than with either modality alone (Wadler S, 2007). These include the incorporation of cetuximab (blocks EGF binding to the EGFR) and the monoclonal antibody bevacizumab (anti VEGF-A) into the various chemotherapy regimes. These combinations show increased response rates and overall survival (Iqbal S *et al*, 2004; Giantonio *et al*, 2006; Kabbinavar FF *et al*, 2006).

## **1.7 Liver Metastases**

### **1.7.1 Summary**

Tumour metastasis is the leading cause of death in cancer patients. The distribution pattern of metastases can be predicted in part by the pattern of regional venous drainage. Most metastases develop in the first capillary bed encountered after discharge from the primary tumour. However, tumours also metastasise to distant locations that are not predictable based on blood flow patterns. The high proportion of bone metastases in breast, prostate and lung cancers are examples of selective homing of tumour cells to a specific organ.

The liver is often the first site of colorectal metastatic disease and may be the only site of spread in as many as 30-40% of patients with advanced disease (Weiss *et al*, 1986). The predilection for metastases to develop in the liver is partly due to the fact that the liver receives the portal drainage from the gastrointestinal tract via the

mesenteric veins. In several experimental tumour models of liver metastasis, metastatic cells first implant in the portal endothelium (Barberaguille E *et al*, 1989). These then break off into a portal radical and re-seed close to the initial lesion. Such satellite formation close to a large liver metastasis is common with secondary tumours of the liver (August DA *et al*, 1985).

For unexplained reasons the right lobe of the liver is more likely to develop metastases than the left lobe (Holbrook RF *et al*, 1995). Macroscopically these metastases are paler in colour than surrounding normal tissue. They enlarge by concentric growth and may attain a large size occupying most of the liver.

Liver metastases may be found in as many as 35% of patients undergoing resection of a colorectal tumour and another 8-30% will subsequently prove to have liver involvement. Furthermore, 40-50% of patients die within 5 years of an apparently curative resection and the liver is the most frequent site of relapse (Scheele J *et al*, 1990; Gordon NLM *et al*, 1993; Stangl R *et al*, 1994; Jatzko *et al*, 1995). 25% of those who develop recurrences have them confined to the liver and therefore may be candidates for surgical resection (Adson MA, 1987). Unfortunately only 5% of all patients with colorectal cancer will ultimately benefit from such intervention (Wagner JS *et al*, 1984). This does, however, translate into 1000 patients treated each year in the UK.

Patients with liver metastases present in the late stages unless detected at the time of the primary cancer or are in a surveillance programme. The symptoms may be weight loss, upper abdominal discomfort and malaise. Evidence of liver failure such as ascites and jaundice are poor prognostic markers.

Inhibition of invasion constitutes a new class of targets for chemo-prevention.

Intervention could commence between the period of tumour proliferation and the

onset of invasion, preventing the series of events leading to metastasis.

### 1.7.2 Detection and Imaging of Liver Metastases

The extent of liver involvement by metastases is an important survival determinant (Wagner JS *et al*, 1984) and, therefore, accurate staging of liver metastases provides a guide to the potential success of surgery. Several staging systems have been proposed (Gennari *et al*, 1986; Gayowski *et al*, 1994). The system modified from the International Union Against Cancer (UICC) and The American Joint Committee on Cancer (AJCC) recommendations for primary hepatobiliary tumours is in popular use. It incorporates tumour size, tumour distribution, number of metastases and extent of extra-hepatic disease (table 1.1).

Stage	5 year survival
I – unilobar solitary tumours of any size	61%
II – unilobar multiple tumours < 2cm without nodal disease	61%
III – unilobar disease with multiple lesions>2cm	28%
IVa – bilobar lesions	20%
IVb – bilobar lesions with nodal disease or extrahepatic disease	0%

Table 1.1 - The staging system modified from the International Union Against Cancer (UICC) and The American Joint Committee on Cancer (AJCC) recommendations for primary hepatobiliary tumours with predicted 5-year survival.

The identification of spread helps to target treatment for those with confined disease and provides valuable prognostic information to all patients with colorectal cancer.

The most widely used imaging technique is a contrast CT scan of the abdomen with

detection rates for hepatic metastases of 68-91% (70% detection for lesions >1cm).

An alternative to the transabdominal ultrasonography approach is intra-operative ultrasound examination (IOUS). IOUS may detect small lesions (<5mm) and other lesions indicating extra hepatic involvement, such as the porta hepatis lymph nodes, which may be difficult to visualise percutaneously (Ravikumar TS, 1996, Kruskal JB *et al*, 1996).

Early experience with laparoscopic ultrasonography suggests it can provide valuable staging information which may alter management (John TJ *et al*, 1994). The combination of a staging laparoscopy and laparoscopic ultrasound improve the selection of candidates for curative resection (Foley EF *et al*, 1998).

A recent meta-analysis of various imaging modalities for detection of colorectal metastases showed that F-18 fluorodeoxyglucose (FDG) PET had significantly higher sensitivity on a per patient basis but not on a per lesion basis compared with other modalities. Sensitivity estimates for MRI imaging with contrast agent were significantly superior to those for helical CT scan with 45 g of iodine or less (Bipat S *et al*, 2005). PET may still miss small hepatic lesions and its performance is affected by recent or current administration of chemotherapy. The list of imaging investigations used for diagnosing liver metastases is outlined in table 1.2. The colon should be fully examined to exclude recurrence or metachronous lesions. Also, the chest needs to be imaged for pulmonary deposits.

### **1.7.3 Current Management of Colorectal Liver Metastases - Surgery**

Patients with untreated liver metastases generally have a poor prognosis, the median survival being less than 12 months (Bengtsson G *et al*, 1981; Lahr CJ *et al*, 1983) and none survive five years. However, survival from time of diagnosis is



Imaging Modality	References
CT	Ward 1989
Transabdominal ultrasound	Ward 1989
Intra-operative ultrasound	Charnley 1991; Hagspiel 1995
Laparoscopic ultrasound	John 1994; Foley 1998
FDG-PET	Bipat 2005
PET / CT / Spiral CT	Votrubova J 2006
MRI	Bipat 2005
Hepatic flow scintigraphy and duplex colour ultrasound (for calculation of hepatic perfusion index)	Leveson 1985; Leen 1991

Table 1.2 – Imaging investigations used in the diagnosis of colorectal liver metastases.

longer in patients with solitary lesions than in those with more widespread involvement (Rougier P *et al*, 1995). Several large series have reported five year survival ranging from 25% to 44% following resection of colorectal liver metastases (Cady B *et al*, 1991; Fong *et al*, 1999; Choti MA *et al*, 1999). A randomised study assessing outcome following resection compared with other treatment modalities is not likely to be undertaken.

Management of all patients with colorectal cancer should be the responsibility of colorectal cancer multidisciplinary teams. More recent data reports lower operative mortality after resection of colorectal liver metastases than that reported in previous decades (Scheele J *et al*, 1995; Stangl R *et al*, 1994; Cady B *et al*, 1991; Choti MA *et al*, 1999; Fong Y *et al*, 1997). No doubt, the reason for this is multifactorial but a volume effect on outcome has been shown by several recent studies (Dimick JB *et al*, 2003).

Resection of hepatic metastases from colorectal cancer has been shown to offer the best chance of long-term survival (Scheele J *et al*, 1995).

Unfortunately the majority of patients with colorectal hepatic metastases are not suitable for resection and therefore specific criteria should be used to select those who may benefit from such intervention. There still remains some controversy amongst clinicians about which patients are most suitable for liver resection and which would be better treated by non-surgical means.

In 1963 Woodington and Waugh published their experience with resection of colorectal liver metastases at the Mayo Clinic and demonstrated that a 20% 5-year survival was possible in selected patients (Woodington GF *et al*, 1963); these results were confirmed by others who reported their experience of liver resection for metastases from a variety of gastrointestinal primary tumours (Flanagan L *et al*, 1967). Although only 7-10% of all patients with colorectal metastases are likely to benefit from hepatic resection, the frequency of this condition means that in the US approximately 5000 patients each year might be suitable for resection of liver metastases (Steele G *et al*, 1991); extrapolated to the UK this suggests that up to 1000 patients each year might be treated in this way with benefit. Attempts to increase resection rates with neo-adjuvant (pre-treatment) chemotherapy have shown promise with a 14% increase in resectability reported in one study (Adam R *et al*, 2001). Neo-adjuvant chemotherapy has been used to convert unresectable tumours to resectable ones using 5-FU, folinic acid and oxaliplatin.

Concerns about the remaining liver reserve after resection have led to the use of preoperative portal vein embolisation to increase the volume of the residual liver (Farges O *et al*, 2003). Another approach has been to plan a two stage



hepatic resection in which the first stage is a non-curative resection or ablation and after subsequent hypertrophy the residual tumour is excised (Adam R *et al*, 2000; Neeleman N *et al*, 1996).

The two most important features of a tumour that influence patient survival in untreated liver metastases are extent of hepatic involvement at the time of diagnosis and the grade of the primary tumour. Multivariate analysis has not shown a consistent result regarding the relevance of the number of lesions on long term survival (Scheele J *et al*, 1995; Fong Y *et al*, 1997; Fong Y *et al*, 1999; Hughes KS *et al*, 1986; Schindl M *et al*, 2005).

It is, therefore, important to define the acceptable residual functioning volume. Acceptable residual liver volume is thought to be a third of the standard liver volume or the equivalent of two liver segments (Garden OJ *et al*, 2006).

Multivariate analysis of various prognostic factors provides a valuable guide to patient selection for hepatic resection and may prevent unnecessary surgery in patients unlikely to derive benefit from the latter (table 1.3).

Tumours that have the capacity to invade and metastasise to lymphatics are likely to be more aggressive than those without this potential. They may conceivably by-pass the liver, by spreading to pre-aortic lymph nodes, and beyond to the thoracic duct and systemic circulation. Although there may be a correlation between primary tumour stage and survival the relationship is not strong enough to preclude liver resection for patients with more advanced primary tumours. Despite this, there is again no consensus view on the correlation between stage of primary tumour and prognosis in published data (Scheele J *et al*, 1995; Nordlinger *et al*, 1992; Jatzko *et al*, 1995; Gayowski TJ *et al*, 1994).

Distribution of metastases	Gayowski 1994; Little 1987
Adequacy of resection margins	Registry of Hepatic Metastases; Hughes 1988; Little 1987
Formal resection or enucleation	Hughes 1988; Scheele 1995; Jatzko 1995, Lise 1990
Age of patient	Fong 1995
Synchronous vs metachronous resection	Hughes 1988; Nordlinger from his book; Lise 1990
Presence of extra-hepatic disease	Rosen 1992; Nordlinger from his book
Tumour stage and grade	Jatzko 1995; Gayowski 1994

Table 1.3 – factors influencing survival following resection of liver metastases

**Survival following resection.** The reported overall 5-year survival rates following hepatic resection of colorectal liver metastases range from 16-50% (Scheele J *et al*, 1995; Jatzko JR *et al*, 1995; Fong Y *et al*, 1999; Ruers & Bleichrodt, 2002)). Studies of the natural history of untreated liver metastases reveal that a minority of patients with limited disease may live for long periods without treatment. Benefit following resection of liver metastases may be gauged best by comparing survival after resection with survival in this group of patients (Wagner JS *et al*, 1984). Even though some of these patients have “favourable” tumour biology, possibly allowing prolonged survival without treatment, long-term survival beyond 5 years is rare without liver resection (Wagner JS *et al*, 1984). This contrasts to patients undergoing hepatic resection where a survival plateau is noted about 5 years after resection (Adson MA, 1987; Steele G *et al*, 1989); the surviving patients subsequently have a life expectancy similar to that of a matched non-cancer cohort (Adson MA, 1987). Repeat resection also seems to provide equal benefit to the first liver resection

(Antoniou A *et al*, 2007; Suzuki *et al*, 2001).

#### 1.7.4 Other Therapies

**Chemotherapy.** Most patients with liver metastases are not suitable for resection and undergo palliative chemotherapy. Others receive adjuvant therapy and a smaller proportion receives neoadjuvant therapy aiming to down-stage unresectable tumours to resectable ones. Liver metastases receive up to 95% of their blood supply via the hepatic artery. Therefore hepatic arterial chemotherapy, through its locoregional effect, produces better response rate in patients with colorectal liver metastases. Some studies showed that this resulted in improved survival (Martin *et al*, 1990) and quality of life (Allen Mersh *et al*, 1994).

It is not possible to cure liver metastases with chemotherapy but the latter can prolong survival and should be considered in all patients not suitable for surgery (Scheithauer W *et al*, 1993; Simmonds PC, 2000; Jonker DJ *et al*, 2000). The chemotherapy may be administered systemically or via the hepatic artery or portal vein. The most commonly used systemic treatment, 5-fluorouracil (5-FU), inhibits thymidylate synthase and results in a tumour response in 5-18% of cases (Rougier P *et al*. In: Nordlinger B 1992). Newer agents such as irinotecan (topoisomerase I inhibitor) have shown improved 1-year survival compared with standard 5-FU regimen (Rougier P *et al*, 1998). The National Institute of Clinical Excellence (NICE) recommends the use of oxaliplatin-based regimens as first line therapy for all patients with non-resectable disease, and irinotecan based regimens for second line therapy after failure of first line treatment (Douillard JY *et al*, 2000; Giacchetti S *et al*, 2000).

Neoadjuvant chemotherapy has been used to convert unresectable tumours to

respectable ones using 5-FU, folinic acid and oxaliplatin. A conversion rate from unresectable to resectable of 38% and a 3-year survival rate of 55% were reported (Bismuth H *et al*, 1996; Giacchetti S *et al*, 1999).

Adjuvant chemotherapy following liver resection uses 5-FU and folinic acid given systemically as a continuous infusion or a bolus. Two randomised studies show no difference in long term survival following hepatic artery infusion (HAI) (Rudroff C *et al*, 1999; Lorenz M *et al*, 1998). A much larger randomised trial comparing adjuvant HAI plus systemic chemotherapy versus adjuvant systemic chemotherapy in a total of 156 patients found that the former group had a significantly higher 2-year disease-free survival (Kemeny N *et al*, 1999).

**Ablative therapies.** The precise role of ablative therapy for hepatic metastases remains unclear. The CLOCC trial should help clarify this issue (<http://www.eortc.be/protoc/Details.asp?Protocol=40004>). For those who decline surgery, have associated co-morbidities that preclude resection or have treatable extrahepatic disease ablative therapy may be indicated. There may also be a role for ablative therapy in patients whose tumours have been downstaged but remain non-resectable (Oshowa A *et al*, 2003; Jansen MC *et al*, 2005; Elias D *et al*, 2003; Solbiati L *et al*, 2001). Resection combined with radiofrequency ablation (RFA) provides a surgical option for those with unresectable liver metastases and may prolong survival (Pawlik TM *et al*, 2003). RFA has also been shown to be effective on its own in the treatment of small liver metastases (Abitabile P *et al*, 2006) with 80% 2 year survival.

## 1.8 Aims

This project aimed to investigate:

1. The effect of ET-1 and its receptor antagonists on colonic cancer cell lines.
  - a. a mechanism for this action was sought .
  - b. ET receptor activation.
2. The ET receptor profile in human colorectal liver metastases tissue.
3. The *in vivo* effect of an oral ET<sub>A</sub> receptor antagonist on CRC liver metastases in an animal model.

## **CHAPTER II**

### **Mechanisms of Endothelin-1 Stimulated Proliferation in Cancer Cells**

## Aims

This chapter investigates the production of ET-1 and mechanisms of ET-1-stimulated proliferation in cancer cell lines. As described in the introduction ET-1 promotes a mitogenic stimulus *in vitro* in a number of cancer cell lines. In the majority of studies the mitogenic signal is transmitted through the ET<sub>A</sub> receptor.

## 2.1 Introduction

Endothelin-1 (ET-1), a 21 amino acid vasoactive peptide (Yanagisawa *et al*, 1988), is a member of a family of three endothelins (Inoue *et al*, 1989). Its actions are mediated via two G protein-coupled receptor subtypes, ET<sub>A</sub> and ET<sub>B</sub> (Arai *et al*, 1990; Sakurai *et al*, 1990). ET-1 stimulates growth of smooth muscle (Komuro *et al*, 1988), fibroblasts (Yamashita *et al*, 1992) and several human cancer cell lines. The latter include ovarian (Bagnato A *et al*, 1999), prostatic (Nelson J *et al*, 1996), meningioma (Kitagawa *et al*, 1994) and CRC (Ali H *et al*, 2000 b) cells. ET<sub>A</sub> receptor antagonists block the growth effect in human ovarian, prostatic and CRC cell lines. The present study investigates mechanisms of ET-1 induced growth and the roles of mitosis and apoptosis. Furthermore, the effect of G-protein inhibition on the ET-1 mitogenic effect was studied.

Two human CRC cell lines and one rat fibrosarcoma cell line were used. The latter was chosen as it was used previously in an animal model of colorectal liver metastases. In the latter model, the role of ET<sub>A</sub> receptor antagonists *in vivo* was investigated and found to reduce tumour growth significantly (Asham E *et al*, 2001).

## **2.2 Materials and Methods**

### **2.2.1 Cell Culture**

Two human colorectal cancer cell lines; SW620 (derived from a metastatic colorectal adenocarcinoma, ECACC, Salisbury, Wiltshire, UK) and LIM1215 (derived from an HNPCC cancer donated by Professor M. O'Hare, Ludwig Institute of Cancer Research, London, UK) and a rat fibrosarcoma cell line MC28 (originally donated by Professor S. Eccles, Institute of Cancer Research, Sutton, Surrey, UK) were used. The cells were routinely cultured at 37 °C in Dulbecco's Modified Eagle Medium (DMEM) containing phenol red pH indicator. The medium was supplemented with 10% foetal calf serum (FCS), L-glutamine (2mM), penicillin (100IU/ml) and streptomycin (100µg/ml) (all reagents bought from Imperial Laboratories, Basingstoke, Hants, UK). The cells were incubated until they reached 80-90% confluence. At this stage the cells were routinely disaggregated with trypsin (1mg/ml in 0.02% EDTA (ethylenediaminetetraacetic acid) in PBS (phosphate buffered saline)), counted and either passaged or used for the experimental studies described below.

### **2.2.2 ET-1 Assays**

On reaching confluence in 75cm<sup>3</sup> flasks, the cells were disaggregated and seeded at a 1:3 split in FCS-containing medium. After 24 hours the cells were incubated in serum free medium (containing glutamine and antibiotics as described above). Supernatants were collected at 24 and 48 hours, spun down to exclude dead cells and debris then stored at -80°C. At the time of supernatant collection the adherent cell numbers were counted using a haemocytometer. ET-1 levels in the media were assayed using an ET-1 kit for an ELISA (Nycomed Amersham, Little Chalfont,



Bucks., UK). The levels were expressed per ml of medium per  $10^6$  cells.

### 2.2.3 Growth Assays

Measurement of cell growth was based on changes in cell number. Responses to the addition of exogenous ET-1 (Sigma Laboratories, Poole, Dorset, UK) were investigated under serum-free (0% FCS) conditions. Cells were plated in 24 well plates at a concentration of  $2.0 \times 10^4$  cells/ml/well in fully supplemented DMEM. After 24 hours the cells were washed (PBS), and incubated in 1ml/well of ET-1 (in concentrations ranging between  $10^{-11}$  to  $10^{-7}$  M) in serum-free (0% FCS) medium. After 48 or 72 hours the cells were washed and growth stopped by the addition of 10% formal saline at 4°C for at least 1 hour.

In the experiments where ET receptor antagonists were used, cells were seeded, as before, for 24 hours. Cells were washed in PBS and then exposed to ET-1 at its most effective dose (determined from the previous experiments for each cell line) as well as one of two ET receptor antagonists; A-127722 (Alexis Corporation, Bingham, Notts., UK), a selective ET<sub>A</sub> receptor antagonist (at concentrations of  $10^{-11}$  to  $10^{-7}$  M) or BQ-788 (Alexis Corporation), a selective ET<sub>B</sub> receptor antagonist (at a concentration of  $10^{-8}$  M), for 48 or 72 hours. Cell growth was stopped by fixation with formal saline as above.

The effect of G-protein inhibition with pertussis toxin (PT) on ET-1 stimulated cell growth was assessed on LIM1215 and SW620 only. Cells were pre-treated for 6 hours with PT then exposed to increasing doses of PT (0.1-100ng/ml)±ET-1 ( $10^{-8}$ M) for 48 hours. Again cells were fixed in formal saline, as above.

Cell number was then determined by the methylene blue assay (Oliver MH *et al*,

1989) and read as absorbance at 650nm. Briefly, the dye chelates in nucleic acids, therefore the larger the amount of nucleic acids (i.e the larger the cell number), the higher the absorbance values.

#### **2.2.4 Immunohistochemistry**

DNA replication (as a measure of proliferation) in cells grown in the presence of ET-1 and/or ET<sub>A</sub> receptor antagonist for 48 or 72 hours in 24 well plates was determined by immunohistochemistry. The protocol followed was described in the previous section, and concentrations used for ET-1 were  $10^{-8}$ M for MC28,  $10^{-9}$ M for LIM1215 and  $10^{-7}$ M for SW620 and the concentrations of A-127722 were  $10^{-11}$ M for MC28,  $10^{-9}$ M for LIM1215 and  $10^{-7}$ M for SW620. The labelling reagent, 5-bromo-2'-deoxyuridine (BrDU), (Dako Ltd., High Wycombe. Bucks, UK) which binds to replicating DNA, was diluted 1:1000 in serum free tissue culture medium. Medium from the cultured cells was replaced with medium containing BrDU and cells were incubated at 37°C for 6 hours. The labelling medium was then removed and the cells washed in phosphate buffered saline (PBS). The cells were fixed in acid-alcohol (90% ethanol, 5% acetic acid, 5% water) at room temperature for at least 30 minutes. The wells were re-hydrated by washing in PBS and then incubated in reconstituted nuclease/anti-5-bromo 2'-deoxyuridine (Dako Ltd.) for 1 hour. After washing, the cells were incubated in peroxidase anti-mouse IgG2a for 30 minutes, washed again and incubated in 3,3'-diaminobenzidine tetrahydrochloride (DAB) for 5 to 10 minutes. The proportion of cells stained with DAB was counted manually (approximately 6 fields per slip), under a microscope, by two independent observers. The stained cells were those in which replicating intracellular DNA had bound BrDU. Our own pilot studies demonstrated that less than 4 hours incubation with BrDU

resulted in a very small proportion of stained cells, while incubations over 10-12 hours resulted in the majority of cells stained. Therefore, optimum exposure was determined at 6-8 hours.

### **2.2.5 Flow Cytometry**

Apoptotic cells were recognised by their ability to bind Annexin V on cell-membrane-exposed phosphatidyl serine. These cells were then identified by flow cytometry.

Cells grown in DMEM and exposed to ET-1 and/or specific endothelin receptor antagonists were harvested with trypsin and washed once in Annexin V binding buffer (BD PharMingen, San Diego, CA, USA). The cells were re-suspended in 1ml HEPES Buffer and 5.5µl of Annexin V-FITC (BD PharMingen) was added. After incubation in the dark at room temperature for 20 minutes, 100µl of a 50µg/ml solution of Propidium Iodide (Sigma) was added to gate out the dead cells.

The cells were analysed by flow cytometry using a FACSCalibur (Becton Dickinson, San Jose, CA, USA) measuring Annexin-FITC fluorescence in the FL1 channel (530/30 bandpass filter) and Propidium Iodide in FL2 (585/42 bandpass filter). 20,000 events were collected. Apoptotic cells were defined as being Annexin positive but Propidium Iodide negative, while late apoptotic/necrotic cells were positive for both Annexin and Propidium iodide.

### **2.2.6 Statistical analysis**

Analysis of all cell culture experiments was carried out using a 1-way ANOVA. If statistically significant, post-hoc analysis was undertaken. A one-way Dunnetts test, set at  $p < 0.05$  was used for ET-1 dose response experiments, and Tukeys honestly significant difference (HSD) test, also set at  $p < 0.05$ , was used for all other analyses.

Means of minimum 6 independent repeats for all experiments were analysed, unless stated otherwise. All graphical data are shown as means and standard deviations (SD). Immunocytochemical data are given as medians  $\pm$  ranges; the Mann-Whitney test was used for analysis with significance set at 0.05.

## **2.3 Results**

### **2.3.1 ET-1 Assay**

The production of ET-1 by the rat fibrosarcoma cell line MC28 and the colorectal cancer cell lines SW620 and LIM1215 were measured in media conditioned for 24 and 48 hours. The normal range for ET-1 is 0.2 to 0.7 fmol/ml cells in human EDTA plasma and the following results were corrected for this. MC28 cells produced ET-1 to the value of 1.1 and 1.3 fmol/ml/ $10^6$  cells at 24 and 48 hours respectively. LIM1215 values were 21.3 and 22.6 fmol/ml/ $10^6$  cells respectively. SW620 ET-1 production measured 11.3 and 8.4 fmol/ml/ $10^6$  cells respectively. The baseline reading for ET-1 amounts in medium only was 0.6 fmol/ml. Results were derived after pooling supernatant and cell numbers from 4 independent repeats. The assay was repeated once and a mean value derived.

### **2.3.2 Growth Assay**

MC28 cells showed an increase in number of cells on addition of ET-1 ( $10^{-7}$  -  $10^{-10}$ M). The biggest effect was seen at an ET-1 concentration of  $10^{-8}$ M with increases of 16.8% and 21.5% at 48 hours and 72 hours respectively (figure 2.1). The selective ET<sub>A</sub> antagonist A127722 by itself did not influence cell growth at any concentration and neither did the selective ET<sub>B</sub> antagonist BQ-788. The stimulatory effect of ET-1 (at the most effective growth-promoting concentration  $10^{-8}$ M) was reversed by the


addition of the ET<sub>A</sub> receptor antagonist, A-127722, at a concentration of 10<sup>-11</sup>M (figure 2.4). No significant change was seen on addition of ET-1 and the ET<sub>B</sub> receptor antagonist, BQ-788 (table 2.1).

LIM1215 cell growth showed similar trends to MC28 cells. The maximal effect of ET-1 was at 10<sup>-9</sup>M (figure 2.2) showing an increase of 11.1% and 14.4% at 48 and 72 hours respectively. Neither antagonist by itself at any concentration had an effect on cell growth. Again the stimulatory effect of ET-1 (10<sup>-9</sup>M) was reversed by A-127722 (figure 2.5). As previously, no effect was seen with BQ788. SW620 cell growth was augmented by ET-1 at concentrations of 10<sup>-7</sup>-10<sup>-11</sup>M. This effect was maximal at 10<sup>-7</sup>M showing increases of 14.1% and 38% at 48 and 72 hours respectively (figure 2.3). A-127722 alone had a dose dependent inhibitory effect in its own right particularly at 10<sup>-7</sup>M (figure 2.7), but BQ788 did not. The stimulatory effect of ET-1 (10<sup>-7</sup>M) on SW620 cell growth was reversed by addition of A-127722 (figure 2.6). The maximal inhibition of growth was by 12.8% and 18.3% at 48 and 72 hours respectively, at an A-127722 concentration of 10<sup>-7</sup> M. BQ788 was ineffective either way.

The mitogenic effect of ET-1 (10<sup>-8</sup>M) at 48 hours was significantly reduced to near control levels on pre-incubation of cells for 6 hours with the selective G-protein antagonist PT. at doses of 1-100ng/ml, PT inhibited ET-1 stimulated growth to near control levels in both cell lines (SW620 and LIM1215). Inhibition was significant with doses of PT above 0.1ng/ml (p<0.01) (figure 2.8).

Figure 2.1

MC28



**Figures 2.1-2.3:** The effect of the addition of endothelin-1 (ET-1) on MC28 (figure 2.1), SW620 (figure 2.2) and LIM1215 (figure 2.3) cell growth at 48 and 72 hours of incubation. The cells were incubated in the presence of ET-1 ( $10^{-11}$  to  $10^{-7}$  M; x axis). Cell growth was measured using the methylene blue assay and read as absorbance (equivalent to cell number) at 650nm. The absolute absorbance values were then converted to ratios (experimental / control; control=1). The results are displayed as mean ratio (S.D) (y axis) and for comparison a line through the control value is shown. Statistically significant growth is shown as ☆  $p < 0.05$  (1-way ANOVA, Dunnetts). The biggest effect on MC28 is seen at ET-1 of  $10^{-8}$  M and this concentration was used for further experiments. For SW620 the biggest effect was seen at ET-1 concentrations of  $10^{-7}$  M, and for LIM1215 it was  $10^{-9}$  M. The data represents the mean (+SD) of 8 independent repeats.

Figure 2.2

SW620






Figure 2.3

LIM1215



**Figure 2.4**

**MC28**



**Figures 2.4-2.6:** MC28 (figure 2.4), SW620 (figure 2.5) and LIM1215 (figure 2.6) cells were incubated in the presence of ET-1 (most effective concentration)  $\pm$  ET<sub>A</sub> (A-127722, A) or ET<sub>B</sub> (not shown) receptor antagonists, or in the presence of ET<sub>A</sub> antagonist (A) alone (x axis). Cell growth was measured using the methylene blue assay and read as absorbance (equivalent to cell number) at 650nm. The absolute absorbance values were then converted to ratios (experimental / control; control=1). The results are displayed as mean ratio (S.D) (y axis) and for comparison a line through the control value is shown. On addition of a combination of ET-1 and ET<sub>A</sub> antagonist the increase in growth is reversed ( ☆  $p < 0.05$  1-way ANOVA, Tukeys HSD) in all three cell lines. For SW620 only (figure 2.5), the addition of the ETA antagonist by itself reduced growth significantly compared to control values ( ☆☆  $p < 0.05$  Mann Whitney test). The concentrations of ET-1 used were  $10^{-8}$ M,  $10^{-7}$ M and  $10^{-9}$ M for MC28, SW620 and LIM1215 respectively. The concentrations of ET<sub>A</sub> antagonist used were  $10^{-11}$ M for MC28 and LIM1215 and  $10^{-7}$ M for SW620.



Figure 2.5

SW620




Figure 2.6

LIM1215






Figure 2.7

SW620



**Figure 2.7:** The effect of the addition of the  $ET_A$  selective antagonist A127722 on SW620 cell growth at 48 and 72 hours of incubation. The cells were incubated in the presence of A127722 ( $10^{-11}$  to  $10^{-7}$ M; x axis). Cell growth was measured using the methylene blue assay and read as absorbance (equivalent to cell number) at 650nm. The absolute absorbance values were then converted to ratios (experimental / control; control=1). The results are displayed as mean ratio (S.D) (y axis) and for comparison a line through the control value is shown. Statistically significant growth is shown as ☆  $p < 0.05$  (1-way ANOVA, Dunnetts). There was dose dependent inhibition, maximal at  $10^{-7}$ M. The data represents the mean (+SD) of 8 independent experiments.

**Figure 2.8**



**Figure 2.8:** Effect of G-protein inhibition with pertussis toxin (PT) on ET-1 stimulated cell growth. Cells were pre-treated for 6hrs with PT, then with increasing doses of PT (0.1-100ng/ml)  $\pm$  ET-1 ( $10^{-8}$ M) for 48hrs. Cell growth was measured using the methylene blue assay and read as absorbance (equivalent to cell number) at 650nm. Data represents the mean  $\pm$  SD of six independent experiments. LIM1215:  $p=0.003$ , SW620:  $p=0.002$  (Tukeys HSD ). ✦, Significant at  $p<0.05$  on post-hoc analysis (Tukeys HSD).

### 2.3.3 Immunohistochemistry

All cell lines showed significant ET-1-induced increases in DNA proliferation. These increases were reversed on further addition of the selective ET<sub>A</sub> antagonist A-

127722.

The proportion of MC28 cells that stained for BrDU (displaying active DNA replication) under control conditions was 51.7% (S.D.  $\pm 3.3$ ). On addition of ET-1 the proportion became 64.3% (S.D.  $\pm 1.3$ ) (a 23% increase,  $p < 0.01$ ) and this returned to approximately baseline levels, 49.4% (S.D.  $\pm 1.1$ ), when the cells were incubated in both ET-1 and A-127722 (table 2.2).

LIM1215 cells followed the same trend. Under control conditions 42.1% (S.D.  $\pm 2.1$ ) of cells stained for BrDU, while 69.7% (S.D.  $\pm 2.1$ ) did so under the influence of ET-1 (67% increase,  $p < 0.01$ ). When ET-1 and A-127722 were added 38.3% (S.D.  $\pm 3.6$ ) of cells stained (table 2.2).

36.5% (S.D.  $\pm 2.4$ ) of SW620 cells stained for BrDU under control conditions. This percentage increased to 43.0% (S.D.  $\pm 0.6$ ) on addition of ET-1 (16% increase,  $p = 0.015$ ) and fell back to 37.4% (S.D.  $\pm 2.3$ ) on addition of both ET-1 and A-127722 (table 2.2).


#### **2.3.4 Flow Cytometry**

Viable cells do not bind Annexin V-BIOTIN or propidium iodide (lower left quadrant). Early apoptotic cells with exposed phosphatidyl serine but intact cell membranes bind Annexin V-BIOTIN but exclude propidium iodide. Fluorescence from this population is displayed in the lower right quadrant. Necrotic or apoptotic cells in terminal stages will be both Annexin V-BIOTIN and propidium iodide positive and are displayed in the upper right quadrant. A small percentage of normal cell death should be expected in routine cultures of untreated cells.


Figure 2.9 displays flow cytograms for MC28 cells. There were no significant changes in the number of cells undergoing apoptosis in the cells exposed to ET-1 (b) as compared to controls (a). Furthermore, the combination of ET-1 and A-127722

**Figure 2.9.** Flow cytograms for MC28 cells. Viable cells are shown in the lower left-hand quadrant of the dot plots (LL). Early apoptotic cell fluorescence is reported in the lower right-hand quadrant (LR). Necrotic or apoptotic cells in terminal stages will be reported in the upper right-hand quadrant (UR).


The cells were incubated in DMEM (a), in the presence of  $10^{-8}$  M ET-1 (b) or in the presence of  $10^{-8}$  M ET-1 and  $10^{-11}$  M ET<sub>A</sub> antagonist, A-127722 (c). The percentage cells in the LR



quadrant were 1.88, 2.27 and 1.71 respectively. This experiment was repeated 10 times for MC28 cells. Similar results were observed in the other cell lines.



C



(c) shows no change either. The results for all cell lines are displayed in table 2.3 and show similar results for LIM1215 and SW620.

## 2.4 Discussion

Endothelin-1 has mitogenic effects on several cancer cell lines including ovarian (Shichiri M *et al*, 1991; Bagnato A *et al*, 1999), prostatic (Nelson *et al*, 1996), meningioma (Kitagawa *et al*, 1994), epithelial carcinoma cell lines derived from human cervix and larynx (Harland S *et al*, 1998) and colorectal (Ali H *et al*, 2000 b). This chapter describes a proliferative effect on colorectal cancer and rat fibrosarcoma cell growth.

The cancer cell lines investigated, all secreted varying concentrations of ET-1 into their medium. The highest concentration was produced by LIM1215 cells and the lowest by MC28 cells. In contrast, the ET-1 plasma level in patients with colorectal cancer has been measured at about 4 pg/ml (Shankar A *et al*, 1998). This represents a 10,000 fold increase compared to ET-1 from cancer cell lines.

However, ET-1 is thought to act in an autocrine or paracrine manner and, therefore, plasma levels do not necessarily give a true indication of action at the cellular level. The fact remains though, that plasma circulating ET-1 levels are in vast excess to those sufficient for cellular action.

*In vitro*, the biggest growth effect of ET-1 was seen on SW620 cells. Interestingly, only SW620 cells reacted to the addition of A-127722 alone. These results may be due to a relatively lower proportion of endogenous ET-1 binding to the receptors in SW620. Thus, the competitive exogenous ET-1 or A-127722 may have a more pronounced effect than on the other cell lines. Alternatively SW620 cells may have a larger density of ET<sub>A</sub> receptors compared to the other two cell lines investigated or that there are different subtypes of the ET<sub>A</sub> receptor with varying response intensities to the receptor antagonists.

The effect of ET-1 is mediated by the ET<sub>A</sub> receptors. This is demonstrated by the reversal of the ET-1 growth effect with the addition of ET<sub>A</sub> receptor antagonists. ET<sub>B</sub> receptor antagonists have no effect. The addition of exogenous ET-1 had a much greater effect on cell growth than endogenous ET-1 since the use of the ET<sub>A</sub> receptor antagonist, A-127722, had no inhibitory effect on endogenous ET-1 except when used on SW620 cells. These observations are supported by work on other cancer cell lines including prostate (Nelson *et al*, 1996) and ovarian (Kitagawa *et al*, 1994). However, there are reasonable suggestions that in both breast cancer (Rajeshkumar *et al*, 2005) and cutaneous melanoma (Bagnato *et al*, 2004) the ET<sub>B</sub> receptor plays a more integral role in mitogenesis than ET<sub>A</sub> receptor.

Changes in cell numbers are a result of the interplay between apoptosis and mitosis. The increase in BrDU uptake suggests that ET-1 stimulates mitogenesis in the cancer cell lines. Radiolabelled thymidine incorporation studies performed by other

workers have also shown an increase in proliferation. Bagnato (1997) working on human ovarian carcinoma cell lines and Kitagawa (1994) working on cultured human meningioma cells both showed dose dependent increases in DNA synthesis in the presence of ET-1. This increase, in both cell lines, was blocked by BQ-123, an ET<sub>A</sub> receptor antagonist. Measuring DNA replication using BrDU offers only a snapshot in time. However, preliminary measurements made after 8 hours of cell incubation with BrDU showed a similar result, while incubation for an hour was insufficient to show staining and 24 hour incubation resulted in the staining of all cells.

The flow cytometry data showed no changes in apoptosis after incubation with ET-1. However, ET-1 induced inhibition of apoptosis has been reported in endothelial cells via ET<sub>B</sub> receptors (Shichiri M *et al*, 1998) and in prostate cancer cells via the ET<sub>A</sub> receptor (Nelson J *et al*, 2005). Eberl and colleagues reported that the mixed endothelin receptor antagonist Bosentan promoted apoptosis in human (HT29 and SW480) and rat (PROb and REGb) colonic carcinoma cells *in vitro* (Eberl LP *et al*, 2000b). The latter study focused on changes detected 24 hours after treatment. In our experience at such a time interval there was no significant increase in net cell number growth. We, therefore, concentrated on findings at 48 hours and longer.

The G-protein inhibitor PT inhibited the mitogenic effect of ET-1 at doses 1-100ng/ml. This suggests that the mitogenic effect in colorectal cancer cells is mediated by PT sensitive G-protein sub-units: Go or Gi. Our findings contrast with results in ovarian cancer cell lines which demonstrated that ET-1 stimulated DNA replication was not inhibited by pre-incubation with PT (Bagnato A *et al*, 1997). The ET-1 mitogenic action in ovarian cancer cells, therefore, appears to be mediated via different G-protein subunits. In addition the mitogenic effect of ET-1 on vascular smooth muscle cells, mediated through the transactivation of the EGFr, requires a



pertussis toxin insensitive G-protein (del Bufalo D *et al*, 2002). However, stimulation of the ET<sub>A</sub> receptor with ET-1 results in receptor coupling with the PT sensitive Go  $\alpha$  and Gi<sub>3</sub>  $\alpha$ -subunits, in addition to the PT insensitive G-proteins Gq/G<sub>11</sub> (Iwasaki H *et al*, 1998).

ET-1 plays a critical role in tumourigenesis and Kim (2005) has shown that enhancement of the  $\beta$ -Catenin signaling pathway activates the ET-1 gene (EDN1). The cell signalling pathways following activation are complex and involve G-protein phosphorylation of multiple pathways including phosphoinositide 3-kinase (PI3K) cascades which trigger calcium channels (Kawanabe Y *et al*, 2002; 2003). ET-1 effects are also mediated partly by protein kinase C (PKC), phospholipase C (PLC) (Sugawara F *et al*, 1996; Bagnato A *et al*, 1997) and phospholipase D (PLD) (Robin P *et al*, 2005). Both of the cyclo-oxygenase -1 and -2 (COX-1 and COX-2) enzymes are involved in tumour progression by promoting tumour angiogenesis, proliferation, survival, invasion and metastasis in several solid tumours. ET-1 has been shown to induce COX-2 and related prostaglandin E<sub>2</sub> release through the activation of the ET<sub>A</sub> receptor (Hughes AK *et al*, 1995). There is also a complex interaction between the mitogen activated protein kinase (MAPK) pathway and ET-1 signaling (Song *et al*, 1996).

The pharmacological blocking of the ET<sub>A</sub> receptor appears to inhibit tumour progression and there is better understanding of the cellular ramifications of such an action.

<b>A. MC28</b>	<b>Control</b>	<b>ET-1</b>	<b>E/BQ</b>
	1	1.24	1.28
	1	1.23	1.25
	1	1.29	1.24
	1	1.21	1.24
	1	1.19	1.26
	1	1.22	1.24
	1	1.21	1.23
	1	1.23	1.24
<b>Mean</b>	<b>1</b>	<b>1.2275</b>	<b>1.2475</b>
<b>SD</b>		<b>0.0296</b>	<b>0.0158</b>

<b>B. SW620</b>	<b>Control</b>	<b>ET-1</b>	<b>E/BQ</b>
	1	1.29	1.23
	1	1.35	1.25
	1	1.33	1.26
	1	1.28	1.26
	1	1.36	1.29
	1	1.33	1.38
	1	1.32	1.41
	1	1.33	1.36
<b>Mean</b>	<b>1</b>	<b>1.32375</b>	<b>1.305</b>
<b>SD</b>		<b>0.0272</b>	<b>0.0682</b>

<b>C. LIM1215</b>	<b>Control</b>	<b>ET-1</b>	<b>E/BQ</b>
	1	1.16	1.21
	1	1.18	1.2
	1	1.18	1.2
	1	1.19	1.19
	1	1.21	1.22
	1	1.22	1.27
	1	1.21	1.26
	1	1.19	1.2
<b>Mean</b>	<b>1</b>	<b>1.1925</b>	<b>1.21875</b>
<b>SD</b>		<b>0.0198</b>	<b>0.0299</b>

**Table 2.1.** The effect of ET-1 and the selective ET<sub>B</sub> receptor antagonist, BQ-788 (BQ) at a concentration of 10<sup>-8</sup>M, on cell growth of MC28 (A) (ET-1 used at 10<sup>-8</sup>M), SW620 (B) (ET-1 at 10<sup>-7</sup>M), and LIM1215 (C) (ET-1 at 10<sup>-9</sup>M) all after 48hrs agent exposure. Cell growth was measured using the methylene blue assay and read as absorbance (equivalent to cell number) at 650nm. The absolute absorbance values were then converted to ratios for ease of presentation (experimental / control; control=1) and the means and standard deviation (SD) calculated. Each result is the sum of 8 repeats.

<b>Cells</b>  <b>Medians</b>  <b>(range)</b>	<b>MC28</b>	<b>SW620</b>	<b>LIM1215</b>
Control	51.7 (+/-3.3)	36.5 (+/-2.4)	42.1 (+/-2.1)
+ ET-1	64.3 (+/-1.3)	43.0(+/-0.6)	69.7 (+/-2.1)
+ ET-1 & A-127722	49.4 (+/-1.1)	37.4 (+/-2.3)	38.3 (+/-3.6)

**Table 2.2.** Percentage of cells positive for BrDU staining after incubation for 72 hours in DMEM alone (control), ET-1 or ET and ET<sub>A</sub> receptor antagonist, A-127722. Cells were visualised by indirect immunohistochemistry using DAB as chromogen and counted manually under the microscope. There was a significant increase in the proportion of MC28 cells binding BrDU on incubation in ET-1 ( $10^{-8}$ M) compared to control ( $p < 0.01$ , Mann Whitney). This was reversed on incubation in ET-1 ( $10^{-8}$ M) and A-127722 ( $10^{-11}$ M) ( $p < 0.01$ ). Similarly, for SW620, incubation in ET-1 ( $10^{-7}$ M) resulted in a significant increase in the percentage of cells binding BrDU ( $p = 0.01$ ) which was reversed on addition of A-127722 ( $10^{-7}$ M) ( $p = 0.015$ ). LIM1215 also bound more BrDU on incubation in ET-1 ( $10^{-9}$ M) ( $p < 0.01$ ) and this increase was reversed on incubation in ET-1 and A-127722 ( $10^{-11}$ M) ( $p < 0.01$ ). Experiments were run at least three times in quadruplicates. The Mann Whitney test was applied to the data throughout.

Cells	MC 28			SW 620		LIM 1215	
	C	E	E/A	E	E/A	E	E/A
repeat 1	1	0.76	0.66	0.46	2.5	0.31	1.09
repeat 2	1	0.76	0.69	0.76	2.48	0.46	1.45
repeat 3	1	0.73	0.99	1.33	1.57	0.43	0.43
repeat 4	1	0.78	0.96	0.8	0.21	0.46	0.51
repeat 5	1	0.37	1.12	0.96	0.25	2.09	1.27
repeat 6	1	0.46	0.74	1.13	1.42	1.65	1
repeat 7	1	1.36	1.1	0.86	1.41	0.74	0.78
repeat 8	1	1.4	1.41	0.74	1.09	1.47	1.21
repeat 9	1	1.23	1.43	0.69	0.3	0.96	1.21
repeat 10	1	1.21	0.91	0.37	0.14		
repeat 11				1.15	1.26		
mean		0.906	1.001	0.841	1.148	0.952	0.994
S.D		0.369	0.272	0.290	0.858	0.639	0.351
p-value		>0.05	>0.05	>0.05	>0.05	>0.05	>0.05

**Table 2.3.** The proportion of apoptotic cells for MC28, SW620 and LIM1215 cells, as visualised by flow cytometry, is displayed. The cells incubated in DMEM only (control, C) were normalised to a value of 1. The proportion of cells exposed to ET-1 (E), and a combination of ET-1 and A-127722 (E/A) were then expressed as a proportion of 1. There were no significant changes in apoptosis in any of the 3 cell lines ( $p > 0.05$  for all experimental limbs, 1-way ANOVA, Tukeys HSD).

## **CHAPTER III**

### **Endothelin Receptor Expression in Colorectal Liver Metastases**

## Aims

This study aims to establish whether or not the ET receptor profile in colorectal liver metastases differs from normal liver tissue and how it compares to previously detected receptor profiles in normal colon and primary colorectal cancer. If there is a detectable change in receptor profile liver metastases may respond differently to ET-1 compared to normal liver cells.

### 3.1 Introduction

Two main subtypes of ET receptor have been cloned from cDNA libraries, ET<sub>A</sub> and ET<sub>B</sub> (Sakurai T *et al*, 1992). ET-1 may act as a growth factor for colorectal as well as other cancer cells via one or both of its receptors, ET<sub>A</sub> and ET<sub>B</sub>.

The development of highly specific agonists and antagonists for these receptors has provided tools for studying the role of ET-1 in tumour vasculature and tumour growth. Radiotracer techniques used in a rat model demonstrated selective tumour vasoconstriction in response to a selective ET<sub>B</sub> receptor agonist. Autoradiography applied to the same model showed regions of intense specific ET<sub>B</sub> receptor binding scattered throughout the tumour mass (Bell KM *et al*, 1998; 1999). Autoradiography has also detected the presence of binding sites (putative receptors) for ET-1 in colorectal cancer specimens and normal colon tissue (Inagaki H *et al*, 1991 a; 1991b; 1992).

Ali (2000a) demonstrated the up-regulation of ET<sub>A</sub> receptors and down-regulation of ET<sub>B</sub> receptors in primary human colorectal cancer tissue. Here, we used *in vitro* receptor autoradiography to study the expression of

endothelin receptor subtypes in human colorectal liver metastases and in adjacent normal liver. Micro-autoradiography has also been used in combination with immunohistochemistry to identify binding to specific cell types in slide-mounted tissues.

## **3.2 Materials and Methods**

### **3.2.1 Tissue collection and preparation**

Following approval from the local ethical committee, patients (n=7) undergoing liver resection for colorectal liver metastases were informed and consented for use of the specimens for our research purposes. At the operation, 1 cm<sup>3</sup> samples of liver metastases (n=13) and adjacent normal liver tissue (n=13) (approximately 5cm away) were excised, and immediately cut into small sections, stored in 1ml cryotubes and preserved frozen in liquid nitrogen.

### **3.2.2 In vitro autoradiography**

ET-1 receptors were identified as described previously (Dashwood M *et al*, 1990). Briefly, 10µm frozen sections were cut on a cryostat, then thaw-mounted onto gelatinised microscope slides and air dried. These were either used immediately for experiments or stored at -70°C for future use.

The slide-mounted sections were incubated in 150pM [<sup>125</sup>I]ET-1 (specific activity 2000 Ci/mM) and non-specific binding was determined by co-incubation of alternate slides in the presence of 500nM unlabelled ET-1. ET receptor subtypes were identified using 150pM [<sup>125</sup>I]-PD151242 for ET<sub>A</sub>

receptors or [ $^{125}$ I]-BQ3020 for ET<sub>B</sub> receptors, specific activity 2000Ci/mM for both (Amersham Pharmacia Biotech, Little Chalfont, Bucks., U.K.).

Autoradiographs generated on 3H Hyperfilm (Amersham Pharmacia Biotech, Little Chalfont, Bucks., U.K.) (figure 1) were semiquantified by densitometric analysis. High-resolution autoradiography was used to identify binding at the cellular level, where slides were coated with a nuclear emulsion (K-2, Ilford, Mobberly, Cheshire, U.K.).

For quantitative assessment images were digitised (Hewlett Packard scanner, Palo Alto, California, USA) and analysed by reading grey densities of the histograms in Adobe Photoshop. In this densitometric analysis, each cm<sup>2</sup> was divided into 255 points, each point given a comparative figure from 0 to 250 (0=black, 250=white), equivalent to disintegrations per minute. Means and standard deviations were calculated.

### **3.2.3 Double labelling**

Radioligand binding to endothelial cells was confirmed on selected tissue sections by using a combination of immunohistochemical and autoradiographic techniques. Here, vascular endothelium was identified using PECAM (platelet endothelial cell adhesion molecule-1, CD31). Slides were incubated in primary antibody, anti-CD31 (diluted 1:200 in PBS pH 7.4, for 30 minutes and indirect immunohistochemistry performed using the Vector alkaline phosphatase method (Vector Laboratories, Peterborough, Cambs, UK), with vector red as the chromogen and counterstained with Mayer's haematoxylin. After tissue processing and immunohistochemical staining, slides were transferred to buffer containing [ $^{125}$ I]-ET-1, [ $^{125}$ I]-PD151242 for



ET<sub>A</sub> receptor or [<sup>125</sup>I]-BQ3020 for ET<sub>B</sub> receptor binding as described previously. Sections were coated with nuclear emulsion and stored for 8 days at 4°C in light-proof boxes containing a dessicant. After processing the emulsion following the manufacturer's instructions [<sup>125</sup>I]-ET-1 binding to endothelial cells was evident as an accumulation of dark grains overlying cells exhibiting positive PECAM immunoreactivity.

### **3.2.4 Statistical analysis**

Analysis of the autoradiography data was carried out using the Wilcoxon signed rank test.

## **3.3 Results**

### **3.3.1 In vitro Autoradiography**

[<sup>125</sup>I]-ET-1 binding was identified in both normal liver and liver metastases sections. Tissue specimens from liver metastases displayed a significant reduction in ET<sub>B</sub> receptor binding ([<sup>125</sup>I]-BQ3020 binding) in tissue from liver metastases ( $64.6 \times 10^3$  gd [grey densities]/mm<sup>2</sup> S.D. 6.9) compared to normal liver tissue ( $134.5 \times 10^3$  gd/mm<sup>2</sup> S.D. 16.9) ( $p < 0.0001$ ). ET<sub>A</sub> receptor density ([<sup>125</sup>I]-PD151242 binding) was not significantly altered in liver metastases ( $103.3 \times 10^3$  gd/mm<sup>2</sup> S.D. 23.2) compared to normal liver tissue ( $116.7 \times 10^3$  gd/mm<sup>2</sup> S.D. 15.6) ( table 3.1 and figure 3.1).

Micro-autoradiographs, generated on emulsion, showed binding of [<sup>125</sup>I]-ET-1 to endothelial cells in the tumour tissue (figure 3.2).

### 3.3.2 Double labelling


[<sup>125</sup>I]-ET-1 binding to vascular endothelial cells within the liver metastases section was confirmed using CD31 (figure 3.3).

Specimen number	Liver metastases		Normal liver	
	ET <sub>A</sub>	ET <sub>B</sub>	ET <sub>A</sub>	ET <sub>B</sub>
	Mean (SD) gdx1000/mm <sup>2</sup>	Mean (SD) gd1000/mm <sup>2</sup>	Mean (SD) gdx1000/mm <sup>2</sup>	Mean (SD) gdx1000/mm <sup>2</sup>
1	94.35 (25.06)	78.55 (14.36)	111.03 (27.84)	126.08 (29.1)
2	88.27 (20.92)	63.84 (11.88)	127.87 (15.8)	154.51 (15.1)
3	85.34 (18.17)	75.85 (18.21)	118.85 (15.21)	151.19 (26.77)
4	91.33 (22.43)	63.67 (25.29)	117.65 (17.32)	111.26 (14.95)
5	90.93 (25.01)	62.52 (21.25)	110.12 (16.17)	111.78 (14.64)
6	124.08 (25.03)	67.86 (16.84)	120.86 (18.4)	138.85 (16.84)
7	123.3 (25.68)	65.8 (8.68)	124.47 (13.52)	133.48 (16.92)
8	76.5 (25.83)	65.21 (15.7)	123.9 (17.99)	159.69 (17.47)
9	76.82 (32.04)	69.73 (18.07)	134 (15.59)	156.41 (15.67)
10	84.19 (23.75)	54.27 (6.11)	144.62 (17.62)	113.28 (19.2)
11	124 (32.81)	56.96 (6.47)	107.12 (23.22)	118.1 (20.82)
12	146.23 (25.57)	57.05 (7.53)	84.62 (19.32)	140.45 (20.45)
13	137.57 (27.92)	58.31 (7.56)	91.81 (21.72)	132.87 (17.19)
Mean	103.3	64.59	116.69	134.46
SD	23.18	6.95	15.58	16.87


**Table 3.1.** Endothelin receptor expression in specimens of human liver metastases and normal liver, displayed in terms of grey density(gd) /mm<sup>2</sup>[equivalent to disintegrations per minute]. The values were derived by semiquantitative densitometry. ET<sub>A</sub> receptor density in liver metastases was not significantly changed (p>0.01 Wilcoxon signed rank test) from that in normal liver (103.30 S.D. 23.18 vs. 116.69 S.D. 15.58 respectively). ET<sub>B</sub> density, however, was significantly (p<0.01 Wilcoxon signed rank test) lower in liver metastases than in normal liver (64.59 S.D. 6.95 vs. 134.46 S.D. 16.87 respectively).

The data is displayed in graph form in **Figure 3.1**


**Figure 3.1**



**Figure 3.2**



$ET_A$  and  $ET_B$  receptor binding on sections of liver metastases demonstrated by gross autoradiography. Semiquantitative densitometric analysis was performed on these films. Total ET binding sites are demonstrated on the left followed by  $ET_A$  then  $ET_B$  binding sites. Non-specific (non spec) binding was excluded from the total.



**Figure 3.3** [125I]-ET-1 binding (dark grain accumulation) to sections of liver metastases using "double labelling". There is dense binding to the endothelial cells (PECAM-positive, red stain) of tumour vessels.

### 3.4 Discussion

ET-1 has been implicated in the pathophysiology of a number of cardiovascular diseases, including coronary syndromes (Salomone O *et al*, 1995) systemic and pulmonary hypertension (Rubanyi GM *et al*, 1994) and atherosclerosis (Dashwood M *et al*, 1993; Zeiher AM *et al*, 1994). Studies using autoradiographic binding studies, demonstrated variations in endothelin receptor expression in human coronary vasculature and varicose veins (Dashwood M *et al*, 1998; Agu O *et al*, 2002). Radioligand binding combined with quantitative autoradiography can measure the affinity and localize the distribution of receptors (Inagaki *et al*, 1991a; 1991b; 1992).

Immunocytochemical localization using receptor subtype selective antisera has also been used successfully to map endothelin receptor distribution (Kuc R *et al*, 2004).

Tumour vasculature is morphologically and functionally abnormal (Carmeliet *et al*, 2000). Therefore, the vascular reactivity differs in tumours and there is potential for therapeutic selective modification of tumour blood flow (Brown *et al*, 1998). The vasoactive action of ET-1 is mediated primarily via the two ET-1 receptors. ET<sub>A</sub> receptors are mainly found on vascular smooth muscle cells and pericytes. These account for the profound vasoconstriction of ET-1. ET<sub>B</sub> receptors are found on contractile elements and endothelial cells and their activation can cause both vasoconstriction and vasodilatation (Davenport AP, 2002).

Additional to vascular pathologies, ET-1 has also been strongly implicated in the pathophysiology of different cancers. It has been shown that ET-1 has mitogenic effects on prostatic (Nelson *et al*, 1996), ovarian (Moraitis *et al*, 1997; Bagnato *et al*, 1999) and colorectal (Ali H *et al*, 2000b) cancer cells in vitro. Shankar (1998) demonstrated that there are elevated levels of ET-1 in the plasma of patients with CRC and colorectal cancer with liver metastases. Working on human pulmonary tumours, Zhao (1995) used a combination of autoradiography and immunocytochemistry to demonstrate that ET-1 binding sites are present on the blood vessels of the tumours. Similar results, using autoradiographic, saturation and competition studies, were obtained by Harland (1998) on human gliomas and meningiomas. In fact the latter group showed that the binding was predominantly to ET<sub>A</sub> receptors. ET<sub>A</sub> receptor expression is up-regulated in several types of cancer cell lines and in human

colorectal cancer tissue. It has been suggested that the mitogenic stimulus of ET-1 is mediated predominantly via the ET<sub>A</sub> receptor (Shichiri M *et al*, 1991; Moraitis S *et al*, 1997). ET-1 stimulates proliferation and migration of endothelial cells through the ET<sub>B</sub> receptor (Morbideilli *et al*, 1995; Wren *et al*, 1993; Goligorsky *et al*, 1999) and is a potent mitogen for vascular smooth muscle and tumour cells through the ET<sub>A</sub> receptor (Alberts *et al*, 1994; Bagnato *et al*, 1998). Nelson (1996), using autoradiography, and Bagnato (1999), using reverse transcription-PCR and Northern blot analysis, both demonstrated ET-1 production and endothelin receptor expression in prostatic and ovarian cancer cells respectively. In both cases an ET<sub>A</sub> receptor antagonist inhibited the ET-1-stimulated growth of cancer cells. Subsequently, Salani and colleagues (2000) localised ET<sub>A</sub> receptor expression to ovarian carcinoma cells and blood vessels whereas ET<sub>B</sub> receptors were confined to vascular endothelial cells. Autoradiographic binding studies confirmed that ovarian carcinoma cells functionally expressed ET<sub>A</sub>, and not ET<sub>B</sub>, receptors and that blood vessels co-expressed both receptors (Salani D *et al*, 2000). Ali (2000a) performed similar work on human CRC cell lines and tissue. This group was able to show, using autoradiography, that both ET<sub>A</sub> and ET<sub>B</sub> receptors were expressed on the CRC cell lines although ET-1 proliferation signals were transmitted only via ET<sub>A</sub> receptors. Furthermore, there was an up-regulation of ET<sub>A</sub> receptors and down-regulation of ET<sub>B</sub> receptors in human CRC tissue. When these results are considered with those showing over-expression of ET-1 in CRC and colorectal liver metastases tissue (Asham E *et al*, 1997b), there appears a strong case for an autocrine / paracrine role for ET-1, acting via the ET<sub>A</sub> receptor, in CRC.

This study further substantiates the work on colorectal cancer tissue (Ali H *et al*, 2000a). ET-1 was identified in both normal liver and liver metastases tissue (unpublished data from our laboratory). Using subtype specific agonists and antagonists, there are demonstrable endothelin receptors in normal and cancerous liver tissue. Therefore, the autocrine / paracrine loop may operate in colorectal liver metastasis. However, there is a reduction in ET<sub>B</sub> receptor density in the cancer tissue compared to controls. This is in agreement with findings in prostate cancer (Nelson J *et al*, 1996). However, in our view, since the metastatic tissue is colonic in origin, one must compare the endothelin receptor densitometry results with those obtained in colorectal cancer and in normal colon.

Ali (2000a) demonstrated that the ratio of ET<sub>A</sub>:ET<sub>B</sub> receptors in normal colon is 0.6:1 (129.2:207.0) and in primary colorectal cancer is 1.7:1 (206.0:122.4). The ratio in colorectal liver metastases is 1.6:1 (103.3:64.6). Thus, there is preservation of the ET<sub>A</sub>:ET<sub>B</sub> density ratio in metastatic tissue from colorectal cancer. The techniques used for the densitometry were identical in both studies. However, very strong conclusions about the significance of absolute densitometric significance can not be drawn unless one has access to cellular micro-autoradiography. Despite this difficulty, it is interesting that apart from the previously reported apparent over-expression of ET<sub>A</sub> receptors in primary CRC, there is a marked under-expression of ET<sub>B</sub> receptors both in primary and metastatic CRC tissue. It is possible that ET-1 contributes to the mitogenic drive stimulating CRC cell growth. The ET<sub>A</sub>:ET<sub>B</sub> ratio increase in both primary and metastatic cancer suggests that an imbalance in this ratio is an event that further drives cancer growth and/or is an established change

“early” in tumourogenesis.

ET<sub>B</sub> receptor agonists have been shown to enhance the suppression of endothelial cell apoptosis by ET-1 (Shichiri M *et al*, 1998). We have already shown in chapter 2 that ET-1 produces its stimulatory effect on human CRC cell lines by potentiation of mitosis which is mediated to some degree by ET<sub>A</sub> receptors. It is plausible that ET<sub>B</sub> receptors also play a role in apoptotic or mitotic changes in CRC. A reduction in ET<sub>B</sub> receptor density may reduce apoptosis or increase mitosis.

Interestingly, it is also revealed that ET-1 binding sites are focused at the endothelial cells of the cancer tissue. Therefore, ET<sub>A</sub> receptors may play a role in neo-angiogenesis in the tumour, thus aiding mitogenesis further, as suggested by our group (Dawas K *et al*, 1999). However, there is evidence from ischaemic human muscle that ET-1 binding sites on microvessels may well be ET<sub>B</sub> receptors (Tsui JC *et al*, 2002).

It has been shown that human primary and metastatic CRC cells display a variation in the expression of endothelin receptors compared to controls. This, combined with the known ET production by CRC *in vivo* and *in vitro*, supports the autocrine / paracrine hypothesis. The ratio of ET<sub>A</sub>:ET<sub>B</sub> receptors changes from normal colon to colorectal cancer, and this change is preserved in liver metastases. The results discussed here raise the possibility of a therapeutic role for endothelin receptor antagonists in the management of colorectal liver metastases.

The next stage was to test this hypothesis in an animal model.



## **CHAPTER IV**

### **The effect of an oral ET<sub>A</sub> receptor antagonist on colorectal liver metastases in an animal model**

## Aims

This study aims to assess the effect of an oral ET<sub>A</sub> receptor antagonist, A-127722, on colorectal liver metastases in an animal model. The rats were inoculated with syngeneic MC28 rat sarcoma cells. These were previously described as having very similar properties to human colorectal cancer cell lines in vitro, in that their growth curves and dose response to ET-1 and ET receptor antagonists are the same (chapter 2).

### 4.1 Introduction

Blood vessels within colorectal liver metastases lack innervation and have an incomplete smooth muscle coat (Ashraf S *et al*, 1996). Thus the tumour blood vessels should not possess any significant neurogenic vasoconstrictor capability. This is useful for manipulation of the proportional delivery of chemotherapeutic agents to the tumour by selective vasoconstriction of normal hepatic vessels. Blood would then shunt into the tumour with a concomitant improvement in drug delivery. An animal model of colorectal liver metastases using syngeneic MC28 sarcoma in Hooded Lister rats is appropriate as a model of human liver metastases. It relies on the fact that the rat tumours have a similar blood vessel anatomical and physiological profile to human tumours, including hypovascularity and expression of ET-1 by the same cell types (Loizidou M *et al*, 1991, Loesch A *et al*, 1997; Ashraf S *et al*, 1997). Colorectal liver metastases develop by haematogenous spread through the portal circulation. A pilot study using the syngeneic rat model of liver metastases demonstrated that the intraportal administration of the ET<sub>A</sub> specific antagonist, BQ-

123, shortly after tumour cells were inoculated via the same route, resulted in significant reduction in tumour load (Asham E *et al*, 2001). These results may be due to the inhibition of the mitogenic effect of ET-1 but there is strong evidence for the anti-angiogenic effect complimenting the former if not instigating it.

## **4.2 Methods**

### **4.2.1 Cell Culture**

MC28 cells were cultured as outlined in chapter 2. Once they reached 80-90 % confluence they were washed twice in PBS, disaggregated using trypsin/EDTA, then resuspended in Hanks medium to obtain a single cell suspension with a final concentration of  $5 \times 10^6$  cells/ml.

### **4.2.2 Animals**

Inbred male Lister Hooded rats weighing 180-300 g were used. The animals were housed in groups of two or three in standard cages and maintained on a routine rat pellet diet. Rats were divided into two groups: experimental (receiving ET<sub>A</sub> antagonist) and control (receiving medium without antagonist). Water (consumption per rat was worked out as roughly 40-45ml per day) was supplemented with the ET<sub>A</sub> antagonist, A-127722 (A-127722 (5mg/kg/day) was diluted [1mg in 1ml medium] in medium composed of 50% of 5% dextrose in a solution of 1M Eq. of NaOH, 30% propylene glycol and 20% ethanol), in the experimental limb and with the medium alone in the control limb. The supplementation was started 3 days prior to inoculation of cancer cells and continued for 17 consecutive days.

Twenty four hours prior to the termination of the experiment BrdU was injected into

all rats intra-peritoneally (3mg/100g body weight). All experiments were carried out under a specific project license in accordance with the UK Animal (Scientific Procedures) Act 1986.

#### **4.2.3 Inoculation of Cancer Cells**

Anaesthesia was induced in the rats using 3% halothane and maintained on 1.5% halothane and 5L/min oxygen administered via a face mask. A midline laparotomy was performed, the caecum and terminal ileum were delivered through the wound and tributaries of the ileocolic vein were identified in the mesentery. With the aid of an operating microscope  $1 \times 10^6$  MC28 cells (0.2 ml.) were injected into the vein using a 27 gauge needle. Pressure was applied at the puncture site on withdrawal of the needle to avoid backflow of blood. After ensuring satisfactory haemostasis the incision was closed using a continuous 3/0 Prolene® suture.

#### **4.2.4 Tissue Specimens**

Terminal anaesthesia was administered to the animals 2 weeks after surgery. The livers were excised, weighed and a note made of the number and size of metastases. The metastases were then dissected out and their total mass measured. The ratio of tumour to total liver mass was calculated (mass of metastases / total mass of liver).

### **4.3 Results**

There were fourteen rats in each limb of the experiment. Three of the rats in the experimental arm and four in the control arm developed mesenteric deposits in the absence of any liver lesions and were excluded. Two of the rats in the experimental

arm and six in the control arm died before the experiment was concluded.

Of the rats that grew liver metastases the T/L ratio in the experimental limb averaged 0.0196 (n=6) whereas in the control limb the ratio was 0.0074 (n=3).

There is no significant difference in the number, size or T/L ratio between the experimental limb and controls (table 4.1).

A-127722	No.	LW	Mets wt	T/L	Control	No.	LW	Mets wt	T/L
1	1	15.03	0.06	0.004	1				
2	25	18.2	1.28	0.07	2	2	11.79	0.05	0.004
3	5	15.14	0.25	0.017	3	11	15.29	0.023	0.0015
4					4				
5					5				
6					6	2	16.87	0.05	0.003
7	1	12.42	0.02	0.002	7				
8					8				
9					9				
10					10				
11	5	14.7	0.18	0.012	11				
12	7	18.6	0.25	0.013	12				
13	4	13.7	0.11	0.008	13	0	19.16	0	
14	18	13.39	1.13	0.084	14	0	17.34	0	

**Table 4.1.** There were 14 rats in each limb of the experiment (A-127722 is the experimental group and the other is a control group). LW = total liver mass in grams, Mets wt = total mass of liver metastases (grams), T/L= ratio of tumour to total liver mass. No = number of metastases in liver.

## 4.4 Discussion

The results of this experiment did not yield much useful information. In this section, amongst other topics, I shall discuss the possible reasons for the unsuccessful outcome.

Efficacy studies in animal tumour models provide an early opportunity to collect preliminary information on toxicity (Arp LH, 1999). When screening and evaluating cytotoxic chemotherapeutic agents, efficacy studies usually include at least 1

dosage level that causes severe toxicity and death. Pathologic evaluation in early efficacy studies may reveal major target organs, dosage/schedule relationships, pharmacokinetic/toxicity relationships, effects of formulation and route of administration, maximum tolerated dose, cause of death, and reversibility of changes in normal tissues. Intra-peritoneal formulations are frequently used to establish proof of concept for promising compounds (hits) from *in vitro* screens; however, these crude formulations may also induce intra-peritoneal inflammation and confound the interpretation of both efficacy and toxicity. Efficacy studies conducted in the later stages of drug discovery may be used to refine the dose and schedule proposed for phase I clinical trials. Efficacy studies in animal tumour models provide useful toxicologic data for screening potential drug candidates, optimising the therapeutic index, and designing both preclinical and clinical development programs.

An animal model using two syngeneic rat tumour lines has already shown that the ET<sub>B</sub> receptor has an important role to play in modification of tumour blood flow. The ET<sub>B</sub> agonist, IRL1620, caused a profound short-term reduction in blood flow in the tumour in the face of an increase in mean arterial blood pressure (Cemazar M *et al*, 2005; Bell *et al*, 1995; 1999). Receptor binding studies also revealed the existence of both ET<sub>A</sub> and ET<sub>B</sub> receptors in the P22 tumour studied and that ET<sub>A</sub> receptors were more abundant than ET<sub>B</sub> receptors.

However, in a rat breast tumour model treatment of animals with ET-1 resulted in increased tumour blood flow compared to normal breast tissue in normal rats. This increase was attenuated by pre treatment with the ET<sub>B</sub> receptor antagonist BQ-788 (Rai A *et al*, 2003). The pilot study that pre dated this experiment showed a significant inhibitory effect of an ET<sub>A</sub> receptor antagonist on the growth of CRC liver

metastases. However, the present larger study does not do the same. There are important differences in the methodologies which will account for this.

A-127722 should be a more potent inhibitor than BQ-123 due to its greater specificity for ET<sub>A</sub> receptors. The concentrations used were both within the quoted ones in the original manufacturer's data sheets. However, the mode of deliveries did differ. The pilot study used an intra-portal route for administration whereas the purpose of this study was to investigate the more amenable oral route and to this end the simplest way of designing this was to add the required dose to a fixed amount of water; the assumption is that rats have a predictable intake of water per day according to weight so that there was minimal variation in the concentrations of the agent between the rats. The direct oro-gastric route of administration, by injecting the agent directly via an oro-gastric tube was considered but not used as the rats soon develop an aversity to this route and administration for 17 days would have been very difficult without anaesthesia. Furthermore, the technique was considered cruel.

It is however, impossible to say what the bio-availability of the oral A-127722 in rats is. Furthermore, some rats were more unwell than others after the inoculation of the MC28 cells under general anaesthesia and during this period the water consumption may have varied significantly for a long period of time. If the agent has an increased efficacy at an early stage of the disease, then that window of time would have been missed while the rat was recovering. There were other concerns with our model. The post operative mortality rate was high as was the incidence of mesenteric deposits in the absence of liver metastases. The mesenteric deposits suggest either inadequate intravascular inoculation or preferential anchoring of the MC28 cell lines used in the mesentery. The MC28 cells might have been cross contaminated; higher passage

numbers were less reliable in their inherent production of endogenous ET-1. The medium used for diluting the A-127722 (dextrose / sodium hydroxide / propylene glycol and ethanol) added further uncertainty in that its effect on the portal circulation is unknown. Several general animal housing issues were also raised around the time of running of this study.

There are other models that have been used and cited. Gervaz (2000) worked on syngeneic BD IX rats and injected DHD K12 colon carcinoma cells into the spleen to produce diffuse liver metastases. Subsequently, TNP-470, a potent angiogenesis inhibitor, was subcutaneously transplanted after tumour implantation on alternate days for 4 weeks. The authors were able to show a significant reduction in the growth of liver metastases. Lauwers (1999), working on 120 WAG/RIJ rats, induced colonic anastomotic tumour and then tested the effect of suramin, an anti angiogenic agent, on intra peritoneal injection. Surprisingly, the tumours grew in size in response to this agent. ET<sub>A</sub> receptor intraperitoneal blockade (Atrasentan) in murine tumour human ovarian carcinoma xenografts grown subcutaneously produced a 65% inhibition in tumour growth (Rosano *et al*, 2004). Moreover it potentiated the effect of the chemotherapeutic agent, cisplatin, significantly.

Anti-angiogenic therapy potentiates radiotherapy in tumour- bearing animals partly because the former reduces hypoxia in the tumour. Furthermore, a combination of anti-angiogenic and cytotoxic therapy has potential for cure in animals with tumours. The effect of the individual agents appears to be merely inhibitory.



## **Chapter V**

### **Summary and Conclusion of Thesis**

### Summary of findings:

1. ET-1 stimulates the growth of colorectal cancer and rat sarcoma cells in vitro. The effect had an optimal dose dependent on the cell line investigated.
2. The stimulant effect of ET-1 is mediated via its ET<sub>A</sub> receptor. Specific blocking of the ET<sub>A</sub> receptor, using A-127722, reversed the ET-1 effect. An ET<sub>B</sub> receptor antagonist did not show any effect.
3. ET-1 increase mitosis as measured by BrDU uptake in the cell lines. Apoptosis was not altered by ET-1.
4. The increase in mitosis is thought to be mediated by a G protein subunit G<sub>o</sub> or G<sub>i</sub> and was blocked by pertussis toxin, a specific blocker.
5. The ratio of ET<sub>A</sub>:ET<sub>B</sub> receptors found in human colorectal liver metastases tissue was increased compared to normal liver tissue taken from the same patients. This correlated positively with findings in colorectal cancer and normal colorectal tissue from previous studies.
6. The oral administration of the ET<sub>A</sub> receptor antagonist, A-127722, in an animal model of colorectal cancer liver metastases did not show any significant outcome change. However, the model may have been inadequate. We used the MC28 cell line which is a sarcoma cell line. The numbers of rats that developed the liver metastases were small and the mode of oral delivery of the drug was not reliable enough. A previous pilot study using the same ET<sub>A</sub> receptor antagonist

administered through an intraportal parenteral route did show a reduction in tumour load. Other animal models have used, amongst many techniques, subcutaneous tumours, intra splenic tumour injections and intra peritoneal drug administration.

Primary colorectal cancer is surgically curable. However, once there is metastasis the 5-year survival becomes much poorer and the median survival is less than 12 months (Bengtsson G et al, 1981; Lahr CJ et al, 1983). Even the selected patients who benefit from resection only achieve 25-25% 5-year survival (Cady B et al, 1991; Fong et al, 1999; Choti MA et al, 1999). There is potential for improving outcome by reducing the incidence of metastases from colorectal primary tumours. ET-1 is a peptide over-expressed by nearly 100% of colonic cancers. A course of an ET<sub>A</sub> antagonist would be combined with resection of the primary tumour. Investigation of the role of ET<sub>A</sub> receptor antagonists in vivo may lead to a reduction in or inhibition of cancer cell metastases.

ET-1 antagonism, as a novel form of solid cancer therapy, may compliment conventional therapy. It has several potential advantages over conventional palliative therapy. In general anti-angiogenic therapy targets a small focus of endothelial cells in capillaries at sites of angiogenesis, bone marrow suppression, gastrointestinal symptoms or hair loss are unlikely side effects (Folkman J, 1995). Several issues need addressing in order to take this work forward:

1. The nature of the intracellular signalling following the activation of the G protein.
2. Does ET-1 expression by colorectal cancer and liver metastases correlate with survival?

3. A better model for testing the efficacy of ET<sub>A</sub> receptor antagonists in animals.
4. What would be the ideal route of administration of the ET<sub>A</sub> receptor antagonist?

The four points outlined above should form the basis of further research into the viability of using ET receptor antagonism as adjuvant treatment for colorectal cancer and liver metastases.

---

## **Bibliography**

- ♦ Abitabile P, Hartl U, Lange J, Maurer CA. Radiofrequency ablation permits an effective treatment for colorectal liver metastasis. *Eur J Surg Oncol* 2007 Feb; 33(1): 67-71.
- ♦ Adam R, Laurent A, Azoulay D, Castaing D, Bismuth H. Two-stage hepatectomy: A planned strategy to treat irresectable liver tumours. *Ann Surg* 2000; 232: 777-85.
- ♦ Adam R, Avisar E, Ariche A *et al.* Five year survival following hepatic resection after neoadjuvant therapy for nonresectable colorectal (liver) metastases. *Ann Surg Oncol* 2001; 8: 347-53.
- ♦ Adson MA. Resection of liver metastases--when is it worthwhile? *World J Surg.* 1987 Aug; 11(4): 511-20.
- ♦ Agu O, Hamilton G, Baker DM, Dashwood MR. Endothelin receptors in the aetiology and pathophysiology of varicose veins. *Eur J Vasc Endovasc Surg* 2002; 23: 165-71.
- ♦ Alafaci C, Jansen I, Arbab MA, Shiokawa Y, Svendgaard NA, Edvinsson L. Enhanced vasoconstrictor effect of endothelin in cerebral arteries from rats with subarachnoid haemorrhage. *Acta Physiol Scand.* 1990 Mar; 138(3): 317-9.
- ♦ Alberts GF, Peifley KA, Johns A, Kleha JF, Winkles JA. Constitutive endothelin-1 over-expression promotes smooth muscle cell proliferation via an external autocrine loop. *J Biol Chem* 1994; 269: 10112-18.
- ♦ Ali H, Dashwood M, Dawas K, Loizidou M, Savage F, Taylor I. Endothelin receptor expression in colorectal cancer. *J. Cardiovasc. Pharmacol.* 2000a; 36(Suppl.1): S69-S71.

- ♦ Ali H, Loizidou M, Dashwood M, Savage F, Sheard C, Taylor I.  
Stimulation of colorectal cancer cell line growth by endothelin-1 and its inhibition by ET<sub>A</sub> antagonists. *Gut* 2000b; 47: 685-688.
- ♦ Allen-Mersh TG, Earlam S, Fordy C, Abrams K, Houghton J. Quality of life and survival with continuous hepatic artery floxuridine infusion for colorectal liver metastases. *Lancet* 1994; 344: 1255-60.
- ♦ Ambiru S, Miyazaki M, Ito H *et al*. Adjuvant regional chemotherapy after hepatic resection for colorectal liver metastases. *Br J Surg* 1999; 86: 1025-31.
- ♦ Antoniou A, Lovegrove RE, Tilney HS, Heriot AG, John TG, Rees M, Tekkis PP, Welsh FK. Meta-analysis of clinical outcome after first and second liver resection for colorectal metastases. *Surgery* 2007 Jan; 141(1): 9-18.
- ♦ Arai H, Hori S, Aramori I, Ohkubo H, Nakanishi S. Cloning and expression of a cDNA encoding an endothelin receptor. *Nature* 1990; 348: 730-2.
- ♦ Arai H, Nakao K, Takaya K, Hosoda K, Ogawa Y, Nakanishi S, Imura H. The human endothelin-B receptor gene. Structural organization and chromosomal assignment. *J Biol Chem*. 1993 Feb 15; 268(5): 3463-70.
- ♦ Arber N, Eagle CJ, Spicak J, Racz I, Dite P, Hajer J, Zavoral M, Lechuga MJ, Gerletti P, Tang J, Rosenstein RB, Macdonald K, Bhadra P, Fowler R, Wittes J, Zauber AG, Solomon SD, Levin B; PreSAP Trial Investigators. Celecoxib for the prevention of colorectal adenomatous polyps. *N Engl J Med* 2006 Aug 31; 355(9): 885-95.

- ♦ Arinami T, Ishikawa M, Inoue A, Yanagisawa M, Masaki T, Yoshida MC, Hamaguchi H. Chromosomal assignments of the human endothelin family genes: the endothelin-1 gene (EDN1) to 6p23-p24, the endothelin-2 gene (EDN2) to 1p34, and the endothelin-3 gene (EDN3) to 20q13.2-q13.3. *Am J Hum Genet* 1991 May; 48(5): 990-6.
- ♦ Arp LH. Tumor models: assessing toxicity in efficacy studies. *Toxicol Pathol.* 1999 Jan-Feb; 27(1): 121-2.
- ♦ Arun C, London NJ, Hemingway DM. Prognostic significance of elevated endothelin-1 levels in patients with colorectal cancer. *Int J Biol Markers.* 2004 Jan-Mar; 19(1): 32-7.
- ♦ Asham E, Loizidou M, Lakhani S, Miller K, Burnstock G, Boulos PB, Taylor I. Expression of endothelin-1 in 98 patients with colorectal cancer. *Eur J Surg Oncol* 1997a; 23: 589.
- ♦ Asham E, Shankar A, Loizidou M, Fredericks S, Miller K, Boulos PB, Burnstock G, Taylor I. A Increased endothelin-1 in colorectal cancer and reduction of tumour growth by ET(A) receptor antagonism. *Br J Cancer.* 2001 Nov 30; 85(11): 1759-63.
- ♦ Ashraf S, Crowe R, Loizidou MC, Turmaine M, Taylor I, Burnstock G. The absence of autonomic perivascular nerves in human colorectal liver metastases. *Br J Cancer.* 1996 Feb; 73(3): 349-59.
- ♦ Ashraf S, Loizidou M, Crowe R, Turmaine M, Taylor I, Burnstock G. Blood vessels in liver metastases from both sarcoma and carcinoma lack perivascular innervation and smooth muscle cells. *Clin Exp Metastasis.* 1997 Sep; 15(5): 484-98.
- ♦ Ashton N. Oxygen and the growth and development of retinal vessels.



Am J Ophthalmol 1966; 62: 412-35.

- ♦ August DA, Sugarbaker PH, Schneider PD. Lymphatic dissemination of hepatic metastases – implications for the follow up and treatment of patients with colorectal cancer. Cancer 1985; 55: 1490-4.
- ♦ Bagnato A, Tecce R, DiCastro V, Catt K. Activation of mitogenic signalling by endothelin-1 in ovarian carcinoma cells. Cancer Res. 1997; 57: 1306-11.
- ♦ Bagnato A, Catt KJ. Endothelins as autocrine regulators of tumor cell growth. Trends Endocrinol Metab 1998; 9: 378-83.
- ♦ Bagnato A, Salani D, Di Castro V, Wu-Wong JR, Tecce R, Nicotra MR, Venuti A, Natali PG. Expression of endothelin-1 and endothelin A receptor in ovarian carcinoma: evidence for an autocrine role in tumour growth. Cancer Res. 1999; 59: 720-7.
- ♦ Bagnato A, Rosano L, Spinella F, Di Castro V, Tecce R, Natali PG. Endothelin B receptor blockade inhibits dynamics of cell interactions and communications in melanoma cell progression. Cancer Res. 2004 Feb 15; 64(4): 1436-43.
- ♦ Barberaguillem E, Alonsovarona A, Vidalvanaclocha F. Selective implantation and growth in rats and mice of experimental liver metastasis in acinar zone-one. Cancer Res 1989; 49: 4003-10.
- ♦ Barst RJ, Langleben D, Badesch D, Frost A, Lawrence EC, Shapiro S, Naeije R, Galie N; STRIDE-2 Study Group. Treatment of pulmonary arterial hypertension with the selective endothelin-A receptor antagonist sitaxsentan. J Am Coll Cardiol. 2006 16; 47(10): 2049-56.

- ♦ Bell KM, Prise VE, Chaplin DJ, Tozer GM. Effects of endothelin-1 and sarafotoxin S6c on blood flow in a rat tumour. *Cardiovasc Pharmacol* 1995; 26(suppl 3): S222-5.
- ♦ Bell KM, Tozer GM, Poole B, Prise VE, Chaolin DJ. Spatial response of the HSN rat tumour to the endothelin B (ETB) receptor agonist IRL1620: comparison to ETB receptor distribution. *Proceedings of the 20<sup>th</sup> European conference on microcirculation Paris, France. International Proceeding Division, Manduzzi Editore 1998, pp 19-23.*
- ♦ Bell KM, Chaplin DJ, Poole BA, Prise VE, Tozer GM. Modification of blood flow in the HSN tumour and normal tissue of the rat by the endothelin ET(B) receptor agonist, IRL1620. *Int J Cancer* 1999; 80: 295-302.
- ♦ Bengtsson G, Carlsson G, Hafstrom L, Jonsson PE. Natural history of patients with untreated liver metastases from colorectal cancer. *Am J Surg* 1981 May; 141(5): 586-9.
- ♦ Bertagnolli MM, Eagle CJ, Zauber AG, Redston M, Solomon SD, Kim K, Tang J, Rosenstein RB, Wittes J, Corle D, Hess TM, Woloj GM, Boissarie F, Anderson WF, Viner JL, Bagheri D, Burn J, Chung DC, Dewar T, Foley TR, Hoffman N, Macrae F, Pruitt RE, Saltzman JR, Salzberg B, Sylwestrowicz T, Gordon GB, Hawk ET; APC Study Investigators. Celecoxib for the prevention of sporadic colorectal adenomas. *N Engl J Med* 2006 Aug 31; 355(9): 873-84.
- ♦ Bhargava S, Stummeyer T, Hotz B, Hines OJ, Reber HA, Buhr HJ, Hotz HG. Selective inhibition of endothelin receptor A as an anti-

angiogenic and anti-proliferative strategy for human pancreatic cancer.

J Gastrointest Surg 2005 May-Jun; 9(5): 703-9.

- ♦ Bipat S, van Leeuwen MS, Comans EF, Pijl ME, Bossuyt PM, Zwinderman AH, Stoker J. Colorectal liver metastases: CT, MR imaging, and PET for diagnosis – meta-analysis. Radiology 2005; 237: 123-31.
- ♦ Bismuth H, Adam R, Levi , Farabos C, Waechter F, Castaing D, Majno P, Engerran L. Resection of nonresectable liver metastases from colorectal cancer following systemic chemotherapy. Ann Surg 1996; 224: 509-22.
- ♦ Bjelke E. Epidemiologic studies of cancer of the stomach, colon and rectum with special emphasis on the role of diets. Vols I-IV. Thesis, University of Minnesota, 1973.
- ♦ Bloch KD, Eddy RL, Shows TB, Quertermous T. cDNA cloning and chromosomal assignment of the gene encoding endothelin 3. J Biol Chem. 1989 Oct 25; 264(30): 18156-61.
- ♦ Bloch KD, Hong CC, Eddy RL, Shows TB, Quertermous T. cDNA cloning and chromosomal assignment of the endothelin 2 gene: vasoactive intestinal contractor peptide is rat endothelin 2. Genomics. 1991 May; 10(1): 236-42.
- ♦ Boarder MR, Marriott DB. Characterization of endothelin-1 stimulation of catecholamine release from adrenal chromaffin cells. J Cardiovasc Pharmacol. 1989; 13 Suppl 5: S223-4.

- ♦ Bobik A, Grooms A, Millar JA, Mitchell A, Grinpukel S. Growth factor activity of endothelin on vascular smooth muscle. *Am J Physiol* 1990 Mar; 258(3 Pt 1): C408-15.
- ♦ Boldrini L, Gisfredi S, Ursino S, Faviana P, Lucchi M, Melfi F, Mussi A, Basolo F, Fontanini G. Expression of endothelin-1 is related to poor prognosis in non-small cell lung carcinoma. *Eur J Cancer*. 2005 Dec; 41(18): 2828-35.
- ♦ Bosari S, Lee AKC, DeLellis RA, Wiley BD, Heatley GJ, Silverman ML. Microvessel quantitation and prognosis in invasive breast carcinoma. *Hum Pathol* 1992; 23: 755-61.
- ♦ Boulanger CM, Luscher TF. Differential effect of cyclic GMP on the release of endothelin-1 from cultured endothelial cells and intact porcine aorta. *J Cardiovasc Pharmacol*. 1991; 17 Suppl 7: S264-6.
- ♦ Breier G, Risau W. The role of VEGF in blood vessel formation. *Trends Cell Biol* 1996; 6: 454-6.
- ♦ Brown JM, Giaccia AJ. The unique physiology of solid tumors: opportunities (and problems) for cancer therapy. *Cancer Res* 1998; 58: 1408-16.
- ♦ Burgess WH, Macaig T. The heparin binding (fibroblast) growth factor family of proteins. *Ann Rev Biochem* 1989; 58: 575-606.
- ♦ Burkitt DP. Epidemiology of cancer of the colon and rectum. 1971. *Dis Colon Rectum*. 1993 Nov; 36(11): 1071-82.
- ♦ Cady B, Stone MD. The role of surgical resection of liver metastases in colorectal carcinoma. *Semin Oncol* 1991; 18: 399-406.

- ♦ Cady B, Stone MD, McDermott WV Jr, Jenkins RL, Bothe A Jr, Lavin PT, Lovett EJ, Steele GD Jr. Technical and biological factors in disease-free survival with hepatic resection for colorectal cancer metastases. *Arch Surg* 1992; 127: 561-8.
- ♦ Cancer Research Campaign. Factsheet 18.1 1993
- ♦ Carducci MA, Padley RJ, Breul J, Vogelzang NJ, Zonnenberg BA, Daliani DD, Schulman CC, Nabulsi AA, Humerickhouse RA, Weinberg MA, Schmitt JL, Nelson JB. Effect of endothelin-A receptor blockade with atrasentan on tumor progression in men with hormone-refractory prostate cancer: a randomized, phase II, placebo-controlled trial. *Clin Oncol*. 2003 Feb 15; 21(4): 679-89.
- ♦ Carmeliet P, Jain RK. Angiogenesis in cancer and other diseases. *Nature* 2000; 407: 249-57.
- ♦ Cemazar M, Wilson I, Prise VE, Bell KM, Hill SA, Tozer GM. The endothelin B (ETB) receptor agonist IRL1620 is highly vasoconstrictive in two syngeneic rat tumour lines: potential for selective tumour blood flow modification. *Br J Cancer* 2005; 93: 98-106.
- ♦ Charnley RM, Morris DL, Dennison AR, Amar SS, Hardcastle JD. Detection of colorectal liver metastases using intraoperative ultrasonography. *Br J Surg*. 1991 Jan; 78(1): 45-8.
- ♦ Choti MA, Bulkley GB. Management of hepatic metastases. *Liver Transpl Surg* 1999; 5: 65-80.
- ♦ Clozel M, Breu V, Gray GA, Kalina B, Loffler BM, Burri K, Cassal JM, Hirth G, Muller M, Neidhart W *et al*. Pharmacological characterization

of bosentan, a new potent orally active nonpeptide endothelin receptor antagonist. *J Pharmacol Exp Ther* 1994 Jul; 270(1): 228-35.

- ♦ Dameron KM, Volpert OV, Tainsky MA, Bouck N. control of angiogenesis in fibroblasts by p53 regulation of thrombospondin-1. *Science* 1994; 265: 1582-4.
- ♦ Dashwood MR, Sykes RM, Collins MJ, Muddle JR, Prehar S, Theodoropoulos S, Yacoub MH. Identification of [<sup>125</sup>I] endothelin binding sites in human coronary tissue. *Neurochem Int* 1990; 18: 439-44.
- ♦ Dashwood MR, Barker SGE, Muddle JR, Yacoub MH, Martin JR. [<sup>125</sup>I]—Endothelin-1 binding to vasa vasorum and regions of neovascularisation in human and porcine blood vessels: a possible for endothelin in intimal hyperplasia and atherosclerosis. *J Cardiovasc Pharmacol* 1993; 22(Suppl 8): S343-6.
- ♦ Dashwood MR, Timm M, Muddle JR, Ong AC, Tippins JR, Parker R, McManus D, Murday AJ, Madden BP, Kaski JC. Regional variations in endothelin-1 and its receptor subtypes in human coronary vasculature: Pathophysiological implications in coronary disease. *Endothelium* 1998; 6(1): 61-70.
- ♦ Davenport AP. International union of pharmacology. XXIX update on endothelin receptor nomenclature. *Pharmacol Rev* 2002; 54: 219-26.
- ♦ Dawas K, Loizidou M, Shankar A, Ali H, Taylor I. Angiogenesis in cancer: the role of endothelin-1. *Annals of the Royal College of Surgeons of England*, 1999; 81(5): 306-10.

- ♦ Del Bufalo D, Di Castro V, Biroccio A, Varmi M, Salani D, Rosano L, Trisciuglio D, Spinella F, Bagnato A. Endothelin-1 protects ovarian carcinoma cells against paclitaxel-induced apoptosis: requirement for Akt activation. *Mol Pharmacol* 2002 Mar; 61(3): 524-32.
- ♦ Dimick JB, Cowan JAJ, Knol JA *et al*. Hepatic resection in the United States: indications, outcomes and hospital procedural volumes from a nationally representative database. *Arch Surg* 2003; 138: 185-91.
- ♦ Donckier J, Michel L, Delos M, Van Beneden R, Havaux X. Endothelin axis expression in medullary thyroid carcinoma: a potential therapeutic target. *Clin Endocrinol (Oxf)*. 2004 Aug; 61(2): 282-4.
- ♦ Douglas SA, Beck GR Jr, Elliott JD, Ohlstein EH. Pharmacological evidence for the presence of three distinct functional endothelin receptor subtypes in the rabbit lateral saphenous vein. *Br J Pharmacol*. 1995 Apr; 114(8): 1529-40.
- ♦ Douillard JY, Cunningham D, Roth AD, Navarro M, James RD, Karasek P, Jandik P, Iveson T, Carmichael J, Alakl M, Gruia G, Awad L, Rougier P. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *Lancet* 2000; 355:1041-7.
- ♦ Eberl LP, Egidy G, Pinet F, Juillerat-Jeanneret L. Endothelin receptor blockade potentiates FasL-induced apoptosis in colon carcinoma cells via the protein kinase C-pathway. *J Cardiovasc Pharmacol* 2000a; 36(5 Suppl 1):S354-6.

- Eberl LP, Valdenaire O, Saintgiorgio V, Jeanin J-F, Juillerat-Jeanneret L. Endothelin receptor blockade potentiates FasL-induced apoptosis in rat colon carcinoma cells. *Int J Cancer* 2000b; 86(2): 182-7.
- Elias D, Ouellet JF, Bellon N, Pignon J-P, Pocard M, Lasser P. Extrahepatic disease does not contraindicate hepatectomy for colorectal liver metastases. *Br J Surg* 2003; 90: 567-74.
- Elshourbagy NA, Korman DR, Wu HL, Sylvester DR, Lee JA, Nuthalaganti P, Bergsma DJ, Kumar CS, Nambi P. Molecular characterization and regulation of the human endothelin receptors. *J Biol Chem*. 1993 Feb 25; 268(6): 3873-9.
- Farges O, Belghiti J, Kianmanesh R, Regimbeau JM, Santoro R, Vilgrain V, Denys A, Sauvanet A. Portal vein embolisation before right hepatectomy: prospective clinical trial. *Ann Surg* 2003; 237: 208-17.
- Ferrara N, Houck K, Jackman L, Leung DW. Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr Rev* 1992; 13: 18-32.
- Flanagan L, Foster JH. Hepatic resection for metastatic cancer. *Am J Surg* 1967 Apr; 113(4): 551-7.
- Foley EF, Kolecki RV, Scirmer BD. The accuracy of laparoscopic ultrasound in the detection of colorectal liver metastases. *Am J Surg* 1998; 176: 262-4.
- Folkman J. Tumor Angiogenesis: Therapeutic implications. *N Engl J Med* 1971; 285: 1182-6.
- Folkman J. Antiangiogenesis: new concept for therapy of solid tumours. *Ann Surg* 1972; 175: 409-16.



- ♦ Folkman J. Clinical applications of research on angiogenesis. *N Engl J Med* 1995; 333(26): 1757-63.
- ♦ Folkman J. Tumour Angiogenesis. In: *The Molecular Basis of Cancer*. Philadelphia: WB Saunders 1995: 206-32.
- ♦ Fong GH, Rossant J, Gertsenstein M, Breitman ML. Role of the flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. *Nature* 1995; 376: 66-70.
- ♦ Fong Y, Blumgart LH, Fortner JG, Brennan MF. Pancreatic or liver resection for malignancy is safe and effective for the elderly. *Ann Surg* 1995; 222: 426-34.
- ♦ Fong Y, Cohen AM, Fortner JG *et al*. Liver resection for colorectal metastases. *J Clin Oncol* 1997; 15: 938-46.
- ♦ Fong Y, Fortner J, Sun RL *et al*. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer : analysis of 1001 consecutive cases. *Ann Surg* 1999; 230: 309-18.
- ♦ Garden OJ, Rees M, Poston GJ, Mirza D, Saunders M, Ledermann J, Primrose JN, Parks RW. Guidelines for resection of colorectal cancer liver metastases. *Gut* 2006; 55(suppl III): iii1-iii8.
- ♦ Gayowski TJ, Iwatsuki S, Madariaga SR *et al*. Experience in hepatic resection for metastatic colorectal cancer – analysis of clinical and pathological risk factors. *Surgery* 1994; 116: 703-11.
- ♦ Gellai M, Jugus M, Fletcher T, Nambi P, Ohlstein EH, Elliott JD, Brooks DP. Nonpeptide endothelin receptor antagonists. V: Prevention and reversal of acute renal failure in the rat by SB 209670. *J Pharmacol Exp Ther*. 1995 Oct; 275(1): 200-6.

- ♦ Gennari L, Doci R, Bignami P, Bozzetti F. Surgical treatment of hepatic metastases from colorectal cancer. *Ann Surg.* 1986 Jan; 203(1): 49-54.
- ♦ Gervaz P, Scholl B, Padrun V, Gillet M. Growth inhibition of liver metastases by the anti-angiogenic drug TNP-470. *Liver* 2000 Apr; 20(2): 108-13.
- ♦ Giacchetti S, Itzhaki M, Gruia G, Adam R, Zidani R, Kunstlinger F, Brienza S, Alafaci E, Bertheault-Cvitkovic F, Jasmin C, Reynes M, Bismuth H, Misset JL, Levi F. Long term survival of patients with unresectable colorectal cancer liver metastases following infusional chemotherapy with 5-fluorouracil, leucovorin, oxaliplatin and surgery. *Ann Oncol* 1999; 10: 663-9.
- ♦ Giacchetti S, Perpoint B, Zidani R, Le Bail N, Faggiuolo R, Focan C, Chollet P, Llory JF, Letourneau Y, Coudert B, Bertheaut-Cvitkovic F, Larregain-Fournier D, Le Rol A, Walter S, Adam R, Misset JL, Levi F. Phase III multicenter randomized trial of oxaliplatin added to chronomodulated fluorouracil-leucovorin as first-line treatment of metastatic colorectal cancer. *J Clin Oncol.* 2000 Jan; 18(1): 136-47.
- ♦ Giantonio BJ, Levy DE, O'Dwyer PJ, Meropol NJ, Catalano PJ, Benson AB III<sup>rd</sup>. A phase II study of high-dose bevacizumab in combination with irinotecan, 5-fluorouracil, leucovorin, as initial therapy for advanced colorectal cancer: results from the Eastern Cooperative Oncology Group study E2200. *Ann Oncol* 2006; 17: 1399-403.
- ♦ Golfman LS, Hata T, Beamish RE, Dhalla NS. Role of endothelin in heart function in health and disease. *Can J Cardiol.* 1993 Sep; 9(7): 635-53.

- ♦ Goligorsky MS, Budzikowski AS, Tsukahara H, Noiri E. Co-operation between endothelin and nitric oxide in promoting endothelial cell migration and angiogenesis. Clin Exp Pharmacol Physiol 1999; 26: 269-71.
- ♦ Gordon NLM, Dawson AA, Bennett B *et al*. Outcome in colorectal adenocarcinoma – 2 7-year studies of a population. Brit Med J 1993; 307: 707-10.
- ♦ Grant K, Knowles J, Dawas K, Burnstock G, Taylor I, Loizidou M. Mechanisms of endothelin-1 stimulated proliferation in colorectal cancer cells. Br J Surg. 2007 Jan; 94(1): 106-12.
- ♦ Hagspiel KD, Neidl KF, Eichenberger AC, Weder W, Marincek B. Detection of liver metastases: comparison of superparamagnetic iron oxide-enhanced and unenhanced MR imaging at 1.5 T with dynamic CT, intraoperative US, and percutaneous US. Radiology 1995; 196: 471-478.
- ♦ Harland S, Kuc R, Pickard J, Davenport A. Expression of endothelin A receptors in human gliomas and meningiomas, with high affinity for the selective antagonist PD156707. Neurosurgery, 1998; 43(4): 890-8.
- ♦ Harris AL. Antiangiogenesis for cancer therapy. Lancet 1997; 349(suppl 11): 13-15.
- ♦ Hashido K, Gamou T, Adachi M, Tabuchi H, Watanabe T, Furuichi Y, Miyamoto C. Truncation of N-terminal extracellular or C-terminal intracellular domains of human ETA receptor abrogated the binding activity to ET-1. Biochem Biophys Res Commun 1992 Sep 30; 187(3): 1241-8.

- ♦ Hawes BE, Luttrell LM, van Biesen T, Lefkowitz RJ.  
Phosphatidylinositol 3-kinase is an early intermediate in the G beta gamma-mediated mitogen-activated protein kinase signaling pathway. J Biol Chem 1996; 271(21): 12133-6.
- ♦ Haynes WG, Davenport AP, Webb DJ. Endothelin: progress in pharmacology and physiology. Trends Pharmacol Sci. 1993 Jun; 14(6): 225-8.
- ♦ Haynes WG, Webb DJ. Contribution of endogenous generation of endothelin-1 to basal vascular tone. Lancet 1994 Sep 24; 344(8926): 852-4.
- ♦ Hickey KA, Rubanyi G, Paul RJ, Highsmith RF. Characterization of a coronary vasoconstrictor produced by cultured endothelial cells. Am J Physiol 1985 May; 248(5 Pt 1): C550-6.
- ♦ Hirata Y, Tikagi Y, Fukuda Y, Marumo F. Endothelin is a potent mitogen for rat vascular smooth muscle cells. Atherosclerosis 1989; 78: 225-8.
- ♦ Holbrook RF, Rodriguezbigas MA, Ramakrishnan K *et al.* Patterns of colorectal liver metastases according to Couinauds segment. Dis Colon Rectum 1995; 38: 245-8.
- ♦ Hori S, Komatsu Y, Shigemoto R, Mizuno N, Nakanishi S. Distinct tissue distribution and cellular localization of two messenger ribonucleic acids encoding different subtypes of rat endothelin receptors. Endocrinology. 1992 Apr; 130(4): 1885-95.
- ♦ Hosoda K, Nakao K, Hiroshi-Arai, Suga S, Ogawa Y, Mukoyama M, Shirakami G, Saito Y, Nakanishi S, Imura H. Cloning and expression of

human endothelin-1 receptor cDNA. FEBS Lett. 1991 Aug 5; 287(1-2): 23-6.

- ♦ Hosoda K, Nakao K, Tamura N, Arai H, Ogawa Y, Suga S, Nakanishi S, Imura H. Organization, structure, chromosomal assignment, and expression of the gene encoding the human endothelin-A receptor. J Biol Chem 1992 Sep 15; 267(26): 18797-804.
- ♦ Huang X, Molema G, King S, Watkins L, Edgington TS, Thorpe PE. Tumour infarction in mice by antibody-directed targeting of tissue factor to tumour vasculature. Science 1997; 275: 547-50.
- ♦ Hughes AK, Padilla E, Kutchera WA, Michael JR, Kohan DE. Endothelin-1 induction of cyclooxygenase-2 expression in rat mesangial cells. Kidney Int 1995 Jan; 47(1): 53-61.
- ♦ Hughes KS, Simon R, Songhorabodi S, Adson MA, Ilstrup DM, Fortner JG, Maclean BJ, Foster JH, Daly JM, Fitzherbert D *et al.* Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of patterns of recurrence. Surgery 1986; 100: 278-84.
- ♦ Hughes KS, Rosenstein RB, Songhorabodi S, Adson MA, Ilstrup DM, Fortner JG, Maclean BJ, Foster JH, Daly JM, Fitzherbert D *et al.* Resection of the liver for colorectal carcinoma metastases. A multi-institutional study of long-term survivors. Dis Colon Rectum 1988; 31: 1-4.
- ♦ Ihara M, Noguchi K, Saeki T, Fukuroda T, Tsuchida S, Kimura S, Fukami T, Ishikawa K, Nishikibe M, Yano M. Biological profiles of highly potent novel endothelin antagonists selective for the ETA receptor. Life Sci 1992; 50(4): 247-55.

- ♦ Imai T, Hirata Y, Emori T, Marumo F. Heparin has an inhibitory effect on endothelin-1 synthesis and release by endothelial cells. *Hypertension* 1993 Mar; 21(3): 353-8.
- ♦ Inagaki H, Bishop AE, Escrig C, Wharton J, Allen-Mersh TG, Polak JM. Localization of endothelin-like immunoreactivity and endothelin binding sites in human colon. *Gastroenterology* 1991a; 101: 47-54.
- ♦ Inagaki H, Bishop AE, Yura J, Polak J. Localization of endothelin-1 and its binding sites in the nervous system of the human colon. *J Cardiovasc Pharmacol* 1991b; 17: S455-7.
- ♦ Inagaki H, Bishop AE, Eimoto , Polak J. Autoradiographic localisation of endothelin-1 binding sites in human colonic cancer tissue. *J Pathol* 1992; 168: 263-7.
- ♦ Inoue A, Yanagisawa M, Kimura S, Kasuya Y, Miyauchi T, Goto K *et al*. The human endothelin family: Three structurally and pharmacologically distinct isopeptides predicted by three separate genes. *Proc Natl Acad Sci USA* 1989; 86(8): 2863-7.
- ♦ Iqbal S, Lenz HJ. Integration of novel agents in the treatment of colorectal cancer. *Cancer Chemother Pharmacol* 2004; 54 Suppl 1: S32-9.
- ♦ Ishikawa K, Ihara M, Noguchi K, Mase T, Mino N, Saeki T, Fukuroda T, Fukami T, Ozaki S, Nagase T, et al. Biochemicand pharmacological profile of a potent and selective endothelin B-receptor antagonist, BQ-788. *Proc Natl Acad Sci U S A*. 1994 May 24; 91(11): 4892-6.
- ♦ Ishikawa T, Yanagisawa M, Kimura S, Goto K, Masaki T. Positive

inotropic action of novel vasoconstrictor peptide endothelin on guinea pig atria. *Am J Physiol* 1988 Oct; 255(4 Pt 2): H970-3.

- ♦ Iwasaki H, Eguchi S, Marumo F, Hirata Y. Endothelin-1 stimulates DNA synthesis of vascular smooth muscle cells through transactivation of epidermal growth factor receptor. *J Cardiovasc Pharmacol* 1998; 31 Suppl 1: S182-4.
- ♦ Jansen MC, van Duijnhoven FH, van Hillegersberg R, Rijken A, van Coevorden F, van der Sijp J, Prevoo W, van Gulik TM. Adverse effects of radiofrequency ablation of liver tumours in the Netherlands. *Br J Surg* 2005; 92: 1248-54.
- ♦ Jatzko GR, Lisborg PH, Stettner HM *et al.* Hepatic resection for metastases from colorectal carcinoma – a survival analysis. *Eur J Cancer* 1995; 31A: 41-6.
- ♦ Jeffery GM, Hickey BE, Hider P. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Library*. Oxford: Update software, 2002; CD002200).
- ♦ John TJ, Greig JD, Crosbie JL, Miles WF, Garden OJ. Superior staging of liver tumors with laparoscopy and laparoscopic ultrasound. *Ann Surg*. 1994 Dec; 220(6): 711-9.
- ♦ Jonker DJ, Maroun JA, Kocha W. Survival benefit of chemotherapy in metastatic colorectal cancer: a meta-analysis of randomised controlled trials. *Br J Cancer* 2000; 82: 1789-94.
- ♦ Kabbinavar FF, Hambleton J, Mass RD, Hurwitz HI, Bergsland E, Sarkar S. Combined analysis of efficacy: the addition of bevacizumab to fluorouracil/leucovorin improves survival for patients with metastatic

colorectal cancer. *J Clin Oncol* 2005 Jun 1; 23(16): 3706-12.

- ♦ Kar S, Yousem S A, Carr B I. Endothelin-1 expression by human hepatocellular carcinoma. *Biochem. Biophys. Res. Commun.* 1995; 216: 514-9.
- ♦ Karaki H, Sudjarwo SA, Hori M, Sakata K, Urade Y, Takai M, Okada T. ETB receptor antagonist, IRL 1038, selectively inhibits the endothelin-induced endothelium-dependent vascular relaxation. *Eur J Pharmacol.* 1993 Feb 16; 231(3): 371-4.
- ♦ Karne S, Jayawickreme CK, Lerner MR. Cloning and characterization of an endothelin-3 specific receptor (ETC receptor) from *Xenopus laevis* dermal melanophores. *J Biol Chem* 1993 Sep 5; 268(25): 19126-33.
- ♦ Kawanabe Y, Hashimoto N, Masaki T. Effects of phosphoinositide 3-kinase on the endothelin-1-induced activation of voltage-independent  $Ca^{2+}$  channels and mitogenesis in Chinese hamster ovary cells stably expressing endothelin A receptor. *Mol Pharmacol* 2002 Sep; 62(3): 756-61.
- ♦ Kawanabe Y, Hashimoto N, Masaki T. Effects of non-selective cation channels and PI3K on endothelin-1-induced PYK2 tyrosine phosphorylation in C6 glioma cells. *Am J Physiol Cell Physiol* 2003 Sep, 285(3), C539-45.
- ♦ Kemeny N, Huang Y, Cohen A, Shi W, Conti JA, Brennan MF, Bertino JR, Turnbull AD, Sullivan D, Stockman J, Blumgart LH, Fong Y. Hepatic arterial infusion of chemotherapy after resection of hepatic metastases from colorectal cancer. *N Engl J Med* 1999; 341: 2039-48.



- ♦ Kikuchi-Yanoshita R, Konishi M, Ito S *et al.* Genetic changes of both p53 alleles associated with the conversion from colorectal adenoma to early carcinoma in familial adenomatous polyposis and non-familial adenomatous polyposis patients. *Cancer Res* 1992; 52: 3965-71.
- ♦ Kim SH, Brennan MF, Russo P *et al.* The role of surgery in the treatment of clinically isolated adrenal metastasis. *Cancer* 1998; 82: 389-94.
- ♦ Kim TH, Xiong H, Zhang Z, Ren B. beta-Catenin activates the growth factor endothelin-1 in colon cancer cells. *Oncogene* 2005 Jan 20; 24(4): 597-604.
- ♦ Kiowski W, Sutsch G, Hunziker P, Muller P, Kim J, Oechslin E, Schmitt R, Jones R, Bertel O. Evidence for endothelin-1-mediated vasoconstriction in severe chronic heart failure. *Lancet* 1995 Sep 16; 346(8977): 732-6.
- ♦ Kishi F, Minami K, Okishima N, Murakami M, Mori S, Yano M, Niwa Y, Nakaya Y, Kido H. Novel 31-amino-acid-length endothelins cause constriction of vascular smooth muscle. *Biochem Biophys Res Commun* 1998 Jul 20; 248(2): 387-90.
- ♦ Kitagawa N, Tsutsumi K, Niwa M, Yamaga SI, Anda T, Khalid H, Himeno A, Taniyama K, Shibata S. A selective endothelin ET<sub>A</sub> antagonist, BQ-123, inhibits <sup>125</sup>I-ET-1 binding to human meningiomas and antagonises ET-1-induced proliferation of meningioma cells. *Cell. Mol. Neurobiol.* 1994; 14(2): 105-118.
- ♦ Kohno M, Yokokawa K, Horio T, Yasunari K, Murakawa K, Takeda T. Atrial and brain natriuretic peptides inhibit the endothelin-1 secretory

response to angiotensin II in porcine aorta. *Circ Res* 1992 Feb; 70(2): 241-7.

- ♦ Kohno M, Horio T, Ikeda M, Yokokawa K, Fukui T, Yasunari K, Murakawa K, Kurihara N, Takeda T. Natriuretic peptides inhibit mesangial cell production of endothelin induced by arginine vasopressin. *Am J Physiol*. 1993 Apr; 264(4 Pt 2): F678-83.
- ♦ Komuro I, Kurihara H, Sugiyama T, Yoshizumi M, Takaku F, Yazaki Y. Endothelin stimulates c-fos and c-myc expression and proliferation of vascular smooth muscle cells. *FEBS Lett* 1988; 238(2): 249-52.
- ♦ Kon V, Sugiura M, Inagami T, Harvie BR, Ichikawa I, Hoover RL. Role of endothelin in cyclosporine-induced glomerular dysfunction. *Kidney Int* 1990 Jun; 37(6): 1487-91.
- ♦ Kooby DA, Fong Y, Suriawinata A *et al*. Impact of steatosis on perioperative outcome following hepatic resection. *J Gastrointest Surg* 2003; 7: 1034-44.
- ♦ Kourembanas S, Marsden PA, McQuillan LP, Faller DV. Hypoxia induces endothelin gene expression and secretion in cultured human endothelium. *J Clin Invest*. 1991 Sep; 88(3): 1054-7.
- ♦ Kronawitter U, Kemeny NE, Heelan R *et al*. Evaluation of chest computed tomography in the staging of patients with potentially resectable liver metastases from colorectal carcinoma. *Cancer* 1999;86: 229-35.
- ♦ Kruskal JB, Kane RA. Imaging of primary and metastatic liver tumors. *Surg Oncol Clin N Am* 1996 Apr; 5(2): 231-60.

- ♦ Kuc R, Davenport A. Comparison of endothelin-A and endothelin-B receptor distribution visualized by radioligand binding versus immunocytochemical localization using subtype selective antisera. *J Cardiovasc Pharmacol* 2004;44(Suppl 1): S224-6.
- ♦ Kurihara H, Yoshizumi M, Sugiyama T, Takaku F, Yanagisawa M, Masaki T, Hamaoki M, Kato H, Yazaki Y. Transforming growth factor-beta stimulates the expression of endothelin mRNA by vascular endothelial cells. *Biochem Biophys Res Commun*. 1989 Mar 31; 159(3): 1435-40.
- ♦ Kusuhara M, Yamaguchi K, Ohnishi A, Abe K, Kimura S, Oona H, Hori S, Nakamura Y. Endothelin potentiates growth factor stimulated DNA synthesis in Swiss 3T3 cells. *Japn J Cancer Res* 1989; 80: 302-5.
- ♦ Kusuhara M, Yamaguchi K, Nagasaki K, Hayashi C, Suzaki A, Hori S, Handa S, Nakamura Y, Abe K. Production of endothelin in human cancer cell lines. *Cancer Res* 1990, 50, 3257-61.
- ♦ Kwak EL, Chung DC. Hereditary colorectal cancer syndromes: an overview. *Clin Colorectal Cancer* 2007 Jan; 6(5): 340-4.
- ♦ Lahr CJ, Soong SJ, Cloud G, Smith JW, Urist MM, Balch CM. A multifactorial analysis of prognostic factors in patients with liver metastases from colorectal carcinoma. *J Clin Oncol* 1983; 1: 720-6.
- ♦ Launois B, Landen S, Heautot JF. Colorectal metastatic liver tumours. In hepatobiliary malignancy. Its multidisciplinary management, ed. J. Terblanche. Arnold, London, 1994, pp. 271-300.
- ♦ Lauwers P, Hubens G, Hendriks J, Vermeulen P, Schuerwegh A, Stevens WJ, De Clerck LS, Dirix L, Van Marck E, Hubens A, Eyskens

E. Effects of suramin on anastomotic colon tumors in a rat model. *Eur Surg Res.* 1999; 31(4): 347-56.

- ♦ Leen E, Goldberg JA, Robertson J, Sutherland GR, Hemingway DM, Cooke TG, McArdle CS. Detection of hepatic metastases using duplex/color Doppler sonography. *Ann Surg.* 1991 Nov; 214(5): 599-604.
- ♦ Lerman A, Edwards BS, Hallet JW, Heublein DM, Sandberg SM, Burnett JC Jr. Circulating and tissue endothelin immunoreactivity in advanced atherosclerosis. *N Engl J Med* 1991, 325, 997-1001.
- ♦ Leveson SH, Wiggins PA, Giles GR, Parkin A, Robinson PJ. Deranged liver blood flow patterns in the detection of liver metastases. *Br J Surg* 1985 Feb; 72(2): 128-30.
- ♦ Lise M, Da Pian PP, Nitti D, Pilati PL, Prevaldi C. Colorectal metastases to the liver: present status of management. *Dis Colon Rectum* 1990 Aug; 33(8): 688-94.
- ♦ Little JM, Hollands M. Hepatic resection for colorectal metastases--selection of cases and determinants of success. *Aust N Z J Surg* 1987 Jun; 57(6): 355-9.
- ♦ Loesch A, Turmaine M, Loizidou M, Crowe R, Ashraf S, Taylor I, Burnstock G. Increase in immunoreactivity for endothelin-1 in blood vessels of rat liver metastases: experimental sarcoma and carcinoma. *J Anat.* 1997 Aug; 191 (Pt 2): 291-9.
- ♦ Loizidou MC, Lawrance RJ, Holt S, Carty NJ, Cooper AJ, Alexander P, Taylor I. Facilitation by partial hepatectomy of tumor growth within the

rat liver following intraportal injection of syngeneic tumor cells. Clin Exp Metastasis 1991; 9(4): 335-49.

- ♦ Lorenz M, Muller HH, Schramm H, Gassel HJ, Rau HG, Ridwelski K, Hauss J, Stieger R, Jauch KW, Bechstein WO, Encke A. Randomised trial of surgery versus surgery followed by adjuvant hepatic arterial infusion with 5-fluorouracil and folinic acid for liver metastases of colorectal cancer. German Cooperative on liver metastases. Ann Surg 1998; 228: 756-62.
- ♦ Marsden PA, Brenner BM. Transcriptional regulation of the endothelin-1 gene by TNF-alpha. Am J Physiol. 1992 Apr; 262(4 Pt 1): C854-61.
- ♦ Martin JK, O'Connell MJ, Wieand HS, Fitzgibbons RJ Jr, Mailliard JA, Rubin J, Nagorney DM, Tschette LK, Krook JE. Intra-arterial floxuridine versus systemic fluorouracil for hepatic metastases from colorectal cancer. Arch Surg 1990; 125:1022-6.
- ♦ Masson S, Latini R, Anand IS, Barlera S, Judd D, Salio M, Perticone F, Perini G, Tognoni G, Cohn JN; Val-HeFT investigators. The prognostic value of big endothelin-1 in more than 2,300 patients with heart failure enrolled in the Valsartan Heart Failure Trial (Val-HeFT). J Card Fail 2006;12(5): 375-80.
- ♦ McClellan G, Weisberg A, Winegrad S. Effect of endothelin-1 on actomyosin ATPase activity. Implications for the efficiency of contraction. Circ Res 1996 Jun; 78(6): 1044-50.
- ♦ Mihara S, Fujimoto M. The endothelin ETA receptor-specific effect of 50-235, a nonpeptide endothelin antagonist. Eur J Pharmacol. 1993 Jun 15; 246(1): 33-8.

- ♦ Millauer B, Longhi MP, Plate KH *et al*. Dominant-negative inhibition of flk-1 suppresses the growth of many tumour types in vivo. *Cancer Res* 1996; 56: 1615-20.
- ♦ Miyauchi T, Yanagisawa M, Tomizausa T, sugishita Y, Suzuki N, Fujino M, Ajisaka R, Goto K, Masaki T. Increased plasma concentrations of endothelin-1 and big endothelin in acute myocardial infarction. *Lancet* 1989; 2(8653): 53-4.
- ♦ Mlynec ML, van Beuningen D, Leder LD, Streffer C. Measurements of the grade of vascularisation in histological tumour tissue sections. *Br J Cancer* 1985; 52: 945-8.
- ♦ Modan B, Barell V, Lubin F, Modan M, Greenberg RA, Graham S. Low-fiber intake as an etiologic factor in cancer of the colon. *J Natl Cancer Inst.* 1975 Jul; 55(1): 15-18.
- ♦ Moraitis S, Langdon SP, Miller WR. Endothelin expression and responsiveness in human ovarian carcinoma cell lines. *Eur. J. Cancer* 1997; 33: 661-8.
- ♦ Morbidelli L, Orlando C, Maggi CA, Ledda F, Ziche M. Proliferation and migration of endothelial cells promoted by endothelins via activation of ET<sub>B</sub> receptors. *Am J Physiol* 1995, 269, 4686-95.
- ♦ Moreland S, McMullen D, Abboa-Offei B, Seymour A. Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature. *Br J Pharmacol.* 1994 Jun; 112(2): 704-8.
- ♦ Morita T, Kurihara H, Maemura K, Yoshizumi M, Yazaki Y. Disruption of cytoskeletal structures mediates shear stress-induced endothelin-1

gene expression in cultured porcine aortic endothelial cells. J Clin Invest. 1993 Oct; 92(4): 1706-12.

- ♦ Mortality Statistics: reviews of the register general on deaths by cause, sex and age in E&W, 1993 (revised 1994) Govt statistical service, 21, 93-4.
- ♦ Nakamuta M, Ohashi M, Tabata S, Tanabe Y, Goto K, Naruse M, Naruse K, Hiroshige K, Nawata H. High plasma concentrations of endothelin like immunoreactives in patients with hepatocellular carcinoma. Am. J. Gastroenterol. 1993; 88(2): 248-52.
- ♦ Neeleman N, Andersson R. Repeated liver resection for recurrent liver cancer. Br J Surg 1996; 83: 893-901.
- ♦ Nelson JB, Hedican SP, George DJ, Reddi AH, Piantodosi S, Eisenberger MA, Simons JW. Identification of endothelin-1 in the pathophysiology of metastatic adenocarcinoma of the prostate. Nat Med 1995; 1(9): 944-9.
- ♦ Nelson J, Cha-Tack K, Hedican S, Magnuson S, Opgenorth T, Bova G, Simons J. Endothelin-1 production and decreased endothelin B receptor expression in advanced prostate cancer. Cancer Res. 1996; 56(4): 663-8.
- ♦ Nelson JB, Carducci MA. The role of the endothelin axis in prostate cancer. Prostate J 1999; 1:126-30
- ♦ Nishimura M, Takahashi H, Matsusawa M, Ikegaki I, Sakamoto M, Nakanishi T, Hirabayashi M, Yoshimura M. Chronic intracerebroventricular infusions of endothelin elevate arterial pressure in rats. J Hypertens. 1991 Jan; 9(1): 71-6.

- Nodan DM in The Development of the Vascular System (eds. Feinberg RN, Sherer GK, Auerbach R) 1-24 Karger, Basel 1991.
- Nordlinger B, Jaeck D, Guiget M, Vaillant JC, Balladur P, Schaal JC. Surgical resection of hepatic metastases: multicentric retrospective study by the French Association of Surgery. In treatment of hepatic metastases of colorectal cancer, ed. B. Nordlinger & D. Jaeck. Springer-Verlag, Paris, 1992, pp. 129-161.
- Oliver MH, Harrison NK, Bishop JE, Cole P, Laurent G. A rapid and convenient assay for counting cells cultured in microwell plates: Application for assessment of growth factors. J. Cell Sci. 1989; 92: 513-8.
- Omland T, Lie RT, Aakvaag A, Aarsland T, Dickstein K. Plasma endothelin determination as a prognostic indicator of 1-year mortality after acute myocardial infarction. Circulation. 1994 Apr; 89(4): 1573-9.
- Opgenorth TJ, Adler AL, Calzadilla SV, Chiou WJ, Dayton BD, Dixon DB, Gehrke LJ, Hernandez L, Magnuson SR, Marsh KC, Novosad EI, Von Geldern TW, Wessale JL, Winn M, Wu-Wong JR. Pharmacological characterization of A-127722: an orally active and highly potent ETA-selective receptor antagonist. J Pharmacol Exp Ther. 1996 Feb; 276(2): 473-81.
- Oshowo A, Gillams AR, Lees W, Taylor I. Radiofrequency ablation extends the scope of surgery in colorectal liver metastases. Eur J Surg Oncol 2003; 29: 244-7.
- Pardanaud L, Yassine F, Dieterlen-Lievre F. Relationship between vasculogenesis, angiogenesis and haematopoiesis during avian



otogeny. Development 1989; 105: 473-85.

- ♦ Patel KV, Schrey MP. Human breast cancer cells contain a phosphoramidon-sensitive metalloproteinase which can process exogenous big endothelin-1 to endothelin-1: a proposed mitogen for human breast fibroblasts. Br J Cancer 1995 Mar; 71(3): 442-7.
- ♦ Pawlik TM, Izzo F, Cohen DS, Morris JS, Curley SA. Combined resection and radiofrequency ablation for advanced hepatic malignancies: results in 172 patients. Ann Surg Oncol 2003; 10: 1059-69.
- ♦ Pipili-Synetos E, Papageorgiou A, Sakkoula E, Sotiropoulo G *et al.* Inhibition of angiogenesis, tumour growth and metastasis by the NO-releasing vasodilators, isosorbide mononitrate and dinitrate. Br J Pharmacol 1995; 116: 1829-34.
- ♦ Potter JD. Colorectal cancer: molecules and populations. J Natl Cancer Inst 1999; 91:916-32.
- ♦ Powell SM, Zilz N, Beazer-Barclay Y, Bryan TM, Hamilton SR, Thibodeau SN, Vogelstein B, Kinzler KW. APC mutations occur early during colorectal tumorigenesis. Nature 1992 Sep 17; 359(6392): 235-7.
- ♦ Rabelink TJ, Kaasjager KA, Boer P, Stroes EG, Braam B, Koomans HA. Effects of endothelin-1 on renal function in humans: implications for physiology and pathophysiology. Kidney Int. 1994 Aug; 46(2): 376-81.

- ♦ Rai A, Gulati A. Evidence for the involvement of ETB receptors in ET-1 induced changes in blood flow to the rat breast tumour. *Cancer Chemother Pharmacol* 2003, 51, 21-8.
- ♦ Rajeshkumar NV, Rai A, Gulati A. Endothelin B receptor agonist, IRL 1620, enhances the anti-tumor efficacy of paclitaxel in breast tumor rats. *Breast Cancer Res Treat.* 2005 Dec; 94(3):237-47.
- ♦ Rakugi H, Tabuchi Y, Nakamaru M, Nagano M, Higashimori K, Mikami H, Ogihara T. Endothelin activates the vascular renin-angiotensin system in rat mesenteric arteries. *Biochem Int.* 1990 Aug; 21(5): 867-72.
- ♦ Ravikumar TS. Laparoscopic staging and intraoperative ultrasonography for liver tumor management. *Surg Oncol Clin N Am* 1996 Apr; 5(2): 271-82.
- ♦ Rees M, Plant G, Wells J *et al.* One hundred and fifty hepatic resections: evolution of technique towards bloodless surgery. *Br J Surg* 1996; 83: 1526-9.
- ♦ Resnick N, Gimbrone MA. Haemodynamic forces are complex regulators of endothelial gene-expression. *FASEB J* 1995; 9: 874-82.
- ♦ Ridge JA, Bading IR, Gelbard AS *et al.* Perfusion of colorectal hepatic metastases – relative distribution of flow from the hepatic artery and portal vein. *Cancer* 1987; 59: 47-53.
- ♦ Riechers H, Albrecht HP, Amberg W, Baumann E, Bernard H, Bohm HJ, Klinge D, Kling A, Muller S, Raschack M, Unger L, Walker N, Wernet W. Discovery and optimization of a novel class of orally active

nonpeptidic endothelin-A receptor antagonists. J Med Chem. 1996 May 24; 39(11): 2123-8.

- ♦ Risau W, Flamme I. Vasculogenesis. Annu Rev Cell Div Biol 1995; 11: 73-91.
- ♦ Risau W. What, if anything, is an angiogenic factor? Cancer Metast Rev 1996; 15: 149-51.
- ♦ Robin P, Chouayekh S, Bole-Feysot C, Leiber D, Tanfin Z.  
Contribution of phospholipase D in endothelin-1-mediated extracellular signal-regulated kinase activation and proliferation in rat uterine leiomyoma cells. Biol Reprod 2005 Jan; 72(1): 69-77.
- ♦ Rosano L, Spinella F, Di Castro V, Natali PG, Bagnato A. Therapeutic targeting of the endothelin-A receptor in human ovarian carcinoma: efficacy of cytotoxic agents is markedly enhanced by co-administration with Atrasentan. J Cardiovasc Pharmacol 2004, 44(suppl 1), S132-5.
- ♦ Rosen CB, Nagorney DM, Taswell HF, Helgeson SL, Ilstrup DM, van Heerden JA, Adson MA. Perioperative blood transfusion and determinants of survival after liver resection for metastatic colorectal carcinoma. Ann Surg. 1992 Oct; 216(4): 493-504.
- ♦ Rougier P, Lasser L, Elias D. Chemotherapy of hepatic metastases of colorectal origin (systemic and local, in palliative or adjuvant treatment). In Nordlinger B, Jaeck D (eds) Treatment of hepatic metastases of colorectal cancer. Paris: Springer-Verlag, 1992, pp. 1092.
- ♦ Rougier P, Laplanche A, Huguier M *et al.* Hepatic arterial infusion of floxuridine in patients with liver metastases from colorectal carcinoma:

long term results of a prospective randomised trial. Clin Oncol 1992; 10: 1112-18.

- ♦ Rougier P, Milan C, Lazorthes F *et al.* Prospective study of prognostic factors in patients with unresected hepatic metastases from colorectal cancer. Br J Surg 1995; 82:1397-400.
- ♦ Rougier P, Van Cutsem E, Bajetta , Niederle N, Possinger K, Labianca R, Navarro M, Morant R, Bleiberg H, Wils J, Awad L, Herait P, Jacques C . Randomised trial of irinotecan versus fluorouracil by continuous infusion after fluorouracil failure in patients with metastatic colorectal cancer. Lancet 1998; 352: 1407-12.
- ♦ Rubanyi GM, Polokoff MA. Endothelins: Molecular biology, biochemistry, pharmacology, physiology and pathophysiology. Pharmacol Rev 1994, 46(3), 325-415.
- ♦ Rudroff C, Altendorf-Hoffmann A, Stagl, Scheele J. Prospective randomised trial on adjuvant hepatic arterial infusion chemotherapy after R0 resection of colorectal liver metastases. Langenbecks Arch Surg 1999, 384,243-9.
- ♦ Ruers T, Bleichrodt RP. Treatment of liver metastases, an update on the possibilities and results. Eur J Cancer 2002; 38(7): 1023-33.
- ♦ Saijonmaa O, Ristimäki A, Fyhrquist F. Atrial natriuretic peptide, nitroglycerine, and nitroprusside reduce basal and stimulated endothelin production from cultured endothelial cells. Biochem Biophys Res Commun. 1990 Dec 14; 173(2): 514-20.

- ♦ Sakurai T, Yanagisawa M, Takuma Y, Miyazaki H, Kimura S, Goto K, Masaki T. Cloning of a cDNA encoding a non-isopeptide selective subtype of the ET receptor. *Nature* 1990; 348: 732-5.
- ♦ Sakurai T, Yanagisawa M, Masaki T. Molecular characterization of endothelin receptors. *Trends Pharmacol Sci.* 1992, 13, 103-8.
- ♦ Sakurai T, Goto K. Endothelins. Vascular actions and clinical implications. *Drugs* 1993 Nov; 46(5): 795-804.
- ♦ Salani D, Di Castro V, Nicotra MR, Rosano L, Tecce R, Venuti A, Natali PG, Bagnato A. Role of endothelin-1 in neovascularisation of ovarian carcinoma. *Am J Path* 2000, 157, 1537-47.
- ♦ Salomone O, Elliott PM, Calvino R, Chester M, Holt D, Kaski JC. Endothelin-1 plasma levels are correlated with the severity of coronary atherosclerosis. *Circulation* 1995; 92(8)Suppl 1: 200.
- ♦ Scheele J, Stangl R, Altendorf-Hofmann A. Hepatic metastases from colorectal carcinoma: impact of surgical resection on the natural history. *Br J Surg* 1990; 77: 1241-6.
- ♦ Scheele J, Stangl R, Altendorf-Hofmann A *et al.* Resection of colorectal liver metastases. *World J Surg* 1995; 19: 59-71.
- ♦ Scheithauer W, Rosen H, Kornek G-V *et al.* Randomised comparison of combination chemotherapy plus supportive care with supportive care alone in patients with metastatic colorectal cancer. *Br Med J* 1993; 306: 2-5.
- ♦ Schindl M, Wigmore SJ, Currie EJ *et al.* Prognostic scoring in colorectal cancer liver metastases: development and validation. *Arch Surg* 2005a; 140: 183-9.

- ♦ Schindl MJ, Redhead DN, Fearon KC *et al.* The value of residual liver volume as a predictor of hepatic dysfunction and infection after major liver resection. *Gut* 2005b; 54: 289-96.
- ♦ Scott N, Bell SM, Sagar P, Blair GE, Dixon MF, Quirke P. p53 expression and K-ras mutation in colorectal adenomas. *Gut* 1993 May; 34(5): 621-4.
- ♦ Scottish Intercollegiate Guidelines Network. 67. Management of colorectal cancer – a national clinical guideline. Edinburgh: Scottish Intercollegiate Guidelines Network, 2003.
- ♦ Shalaby F, Rossant J, Yamaguchi TP, Gertsenstein M, Wu XF, Breitman ML, Schuh AC. Failure of blood island formation and vasculogenesis in flk-1 deficient mice. *Nature* 1995; 376: 62-66.
- ♦ Shankar A, Loizidou M, Aliev G, Fredericks S, Holt D, Boulos PB, Burnstock G, Taylor I. Elevated endothelin-1 production and secretion by cancer. *Br J Surg* 1996; 83: 1623-1652.
- ♦ Shankar A, Loizidou M, Aliev G, Fredericks S, Holt D, Boulos PB, Burnstock G, Taylor I. Raised endothelin-1 levels in patients with colorectal liver metastases. *Br. J. Surg.* 1998; 85: 502-6.
- ♦ Shaw IM, Rees M, Welsh FKS *et al.* Repeat hepatic resection for recurrent colorectal liver metastases is associated with favourable long term survival. *Br J Surg* 2006; 93: 457-64.
- ♦ Shichiri M, Hirata Y, Nakajima T, Ando K, Imai T, Yanagisawa M, Masaki T, Marumo F. Endothelin-1 is an autocrine/paracrine growth factor for human cancer cell lines. *J. Clin. Invest.* 1991; 87(5): 1867-71.

- ♦ Shichiri M, Marumo F, Hirata Y. Endothelin B receptor-mediated suppression of endothelial apoptosis. *J Cardiovasc Pharmacol.* 1998; 31(Suppl.1) : S138-41.
- ♦ Shirouzu K, Isomoto H, Hayashi A et al. Surgical treatment for patients with pulmonary metastases after resection of primary colorectal carcinoma. *Cancer* 1995; 76: 393-8.
- ♦ Short RHD. Alveolar epithelium in relation to growth of the lung. *Phil Trans R Soc Lond* 1950; B235: 35-87.
- ♦ Shoup M, Gonen M, D'Angelico M *et al.* Volumetric analysis predicts hepatic dysfunction in patients undergoing major liver resection. *J Gastrointest Surg* 2003; 7: 325-30.
- ♦ Simmonds PC. Palliative chemotherapy for advanced colorectal cancer: systematic review and meta-analysis. *Colorectal Cancer Collaborative Group. BMJ* 2000; 321: 531-5.
- ♦ Simonson MS, Herman WH. Protein kinase C and protein tyrosine kinase activity contribute to mitogenic signalling by endothelin-1. Cross-talk between G protein-coupled receptors and pp60c-src. *J Biol Chem* 1993; 268(13): 9347-57.
- ♦ Solbiati L, Livraghi T, Goldberg SN, Ierace T, Meloni F, Dellanoce M, Cova L, Halpern EF, Gazelle GS. Percutaneous radio-frequency ablation of hepatic metastases from colorectal cancer: long term results in 117 patients. *Radiology* 2001; 221: 159-66.
- ♦ Song HJ, Min YS, Shin CY, Jeong JH, Sohn UD. Activation of p38 MAPK is involved in endothelin-1-stimulated COX-2 expression in cultured Feline esophageal smooth muscle cells. *Mol Cells* 2006 Aug

- 31; 22(1): 44-50.
- ♦ Spinella MJ, Malik AB, Everitt J, Andersen TT. Design and synthesis of a specific endothelin 1 antagonist: effects on pulmonary vasoconstriction. *Proc Natl Acad Sci U S A*. 1991 Aug 15; 88(16): 7443-6.
  - ♦ Stangl R, Altendorf-Hofmann A, Charnley RM *et al*. Factors influencing the natural history of colorectal liver metastases. *Lancet* 1994;343: 1405-10.
  - ♦ Steele G, Bleday R, Mayer RJ, Lindblad A, Petrelli N, Weaver D. A prospective evaluation of hepatic resection for colorectal carcinoma metastases to the liver: Gastrointestinal Tumor Study Group Protocol 6584. *J Clin Oncol*. 1991 Jul; 9(7): 1105-12.
  - ♦ Steele G, Ravikumar TS. Resection of hepatic metastases from colorectal cancer. Biologic perspective. *Ann Surg*. 1989 Aug; 210(2):127-38.
  - ♦ Stewart GD, O'Suilleabhain CB, Madhavan KK *et al*. The extent of resection influences outcome following hepatectomy for colorectal liver metastases. *Eur J Surg Oncol* 2004; 30: 370-6.
  - ♦ Sudjarwo SA, Hori M, Takai M, Urade Y, Okada T, Karaki H. A novel subtype of endothelin B receptor mediating contraction in swine pulmonary vein. *Life Sci*. 1993; 53(5): 431-7.
  - ♦ Sugawara F, Ninomiya H, Okamoto Y, Miwa S, Mazda O, Katsura Y, Masaki T. Endothelin-1 induced mitogenic responses of Chinese hamster ovary cells expressing human endothelin A: the role of a



wortmannin-sensitive signaling pathway. *Mol Pharmacol* 1996 Mar, 49(3), 447-57.

- ♦ Suzuki S, Sakaguchi T, Yokoi Y, Kurachi K, Okamoto K, Okumura T, Tsuchiya Y, Nakamura T, Konno H, Baba S, Nakamura S. Impact of repeat hepatectomy on recurrent colorectal liver metastases. *Surgery* 200;129(4) :421-8.
- ♦ Tabuchi Y, Nakamaru M, Rakugi H, Nagano M, Mikami H, Ogihara T. Endothelin inhibits presynaptic adrenergic neurotransmission in rat mesenteric artery. *Biochem Biophys Res Commun.* 1989 Jun 15; 161(2): 803-8.
- ♦ Takebayashi Y, Akiyama S-I, Yamada K, Akiba S, Aikou T. Angiogenesis is an unfavourable prognostic factor in human colorectal carcinoma. *Cancer* 1996; 78(2): 226-31.
- ♦ Teicher BA, Holden SA, Ara G *et al.* Potentiation of cytotoxic cancer therapy by TNP-470 alone and with other anti-angiogenic agents. *Int J Cancer* 1994; 57: 920-5.
- ♦ Topal B, Kaufman L, Aerts R *et al.* Patterns of failure following curative resection of colorectal liver metastases. *Eur J Surg Oncol* 2003;29: 248-53.
- ♦ Tsui JC, Baker DM, Biecker E, Shaw S, Dashwood MR. Potential role of endothelin 1 in ischaemia-induced angiogenesis in critical leg ischaemia. *Br J Surg* 2002; 89(6):741-7.
- ♦ Tsujii M, Kawano S, Tsuji S, Sawaoka H, Hori M, DuBois RN. Cyclooxygenase regulates angiogenesis by colon cancer cells. *Cell* 1998; 93: 705-16.

- ♦ Vacca F, Bagnato A, Catt KJ, Tecce R. Transactivation of the epidermal growth factor receptor in endothelin-1-induced mitogenic signaling in human ovarian carcinoma cells. *Cancer Res* 2000, 60(18), 5310-7.
- ♦ Vasen HFA, Watson P, Jukka-Pekka M, Lynch HT and the ICG-HNPCC. New clinical criteria for HNPCC Proposed by the International Collaborative Group on HNPCC. *Gastroenterol* 1999; 116: 1453-6.
- ♦ Vibert E, Perniceni T, Levard H et al. Effect of surgical margin status on survival and site of recurrence after hepatic resection for colorectal metastases. *Br J Surg* 2006; 93: 67-72.
- ♦ Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, Leppert M, Nakamura Y, White R, Smits AM, Bos JL. Genetic alterations during colorectal-tumor development. *N Engl J Med*. 1988 Sep 1; 319(9): 525-32.
- ♦ Votrubova J, Belohlavek O, Jaruskova M, Oliverius M, Lohynska R, Trskova K, Sedlackova E, Lipska L, Stahalova V. The role of FDG-PET/CT in the detection of recurrent colorectal cancer. *Eur J Nucl Med Mol Imaging*. 2006 Jul; 33(7): 779-84.
- ♦ Vukasin AP, Ballantyne GH, Flannery JT, Lerner E, Modlin IM. Increasing incidence of cecal and sigmoid carcinoma. Data from the Connecticut Tumor Registry. *Cancer* 1990 Dec 1 ; 66(11): 2442-9.
- ♦ Wadler S. Targeted therapy in colorectal cancer. *Clin Colorectal Cancer* 2007 Jan; 6(5): 357-61.
- ♦ Wagner JS, Adson MA, Van Heerden JA, Adson MH, Ilstrup DM. The natural history of hepatic metastases from colorectal cancer. A

comparison with resective treatment. *Ann Surg.* 1984 May; 199(5): 502-8.

- ♦ Wagner OF, Christ G, Wojta J, Vierhapper H, Parzer S, Nowotny PJ, Schneider B, Waldhausl W, Binder BR. Polar secretion of endothelin-1 by cultured endothelial cells. *J Biol Chem.* 1992 Aug 15; 267(23): 16066-8.
- ♦ Wanebo HJ, Chu QD, Avradopoulos KA, *et al.* Current perspectives on repeat hepatic resection for colorectal carcinoma: a review. *Surgery* 1996; 119: 361-71.
- ♦ Ward BA, Miller DI, Frank JA *et al.* Prospective evaluation of hepatic imaging studies in the detection of colorectal metastases: correlation with surgical findings. *Surgery* 1989; 105: 180-7.
- ♦ Warner TD, Allcock GH, Corder R, Vane JR. Use of the endothelin antagonists BQ-123 and PD 142893 to reveal three endothelin receptors mediating smooth muscle contraction and the release of EDRF. *Br J Pharmacol.* 1993a Oct; 110(2): 777-82.
- ♦ Warner TD, Allcock GH, Mickley EJ, Corder R, Vane JR. Comparative studies with the endothelin receptor antagonists BQ-123 and PD 142893 indicate at least three endothelin receptors. *J Cardiovasc Pharmacol* 1993b; 22 Suppl 8: S117-20.
- ♦ Watanabe T, Suzuki N, Shimamoto N, Fujino M, Imada A. Endothelin in myocardial infarction. *Nature* 1990, 344, 114.
- ♦ Wei C-M, Lerman A, Rodeheffer RJ, McGregor CG, Brandt RR, Wright S, Heublein DM, Kao PC, Edwards WD, Burnett JC Jr. Endothelin in human congestive heart failure. *Circulation.* 1994 Apr; 89(4): 1580-6.

- ♦ Weidner N, Semple JP, Welch WR, Folkman J. Tumour angiogenesis and metastasis-correlation in invasive breast carcinoma. *N Engl J Med* 1991; 324: 1-8.
- ♦ Weidner N. Intratumour microvessel density as a prognostic factor in cancer. *Am J Pathol* 1995; 147: 9-19.
- ♦ Weiss L, Grundmann E, Torhorst J *et al.* Haematogenous metastatic patterns in colonic carcinoma: an analysis of 1541 necropsies. *J Pathol* 1986; 150: 195-203.
- ♦ Wigmore SJ, Redhead DN, Yan XJ *et al.* Virtual hepatic resection using three-dimensional reconstruction of helical computed tomography angioportograms. *Ann Surg* 2001; 233: 221-6.
- ♦ Wong-Dusting HK, La M, Rand MJ. Mechanisms of the effects of endothelin on responses to noradrenaline and sympathetic nerve stimulation. *Clin Exp Pharmacol Physiol.* 1990 Apr; 17(4): 269-73.
- ♦ Woodington GF, Waugh JM. Results of resection of metastatic tumors of the liver. *Am J Surg* 1963; 105: 24-9.
- ♦ Wren AD, Hiley CR, Fan TD. Endothelin-3 mediated proliferation in wounded human umbilical vein endothelial cells. *Biochem Biophys Res Comm* 1993, 196, 369-75.
- ♦ Wu JS, Paul P, McGannon EA, Church JM. APC genotype, polyp number, and surgical options in familial adenomatous polyposis. *Ann Surg* 1998; 227: 57-62.
- ♦ Wulfig P, Diallo R, Kersting C, Wulfig C, Poremba C, Greb RR, Bocker W, Kiesel L. Endothelin-1, Endothelin-A- and Endothelin-B-

- receptor expression in preinvasive and invasive breast disease. *Oncol Rep.* 2004 Apr; 11(4): 791-6.
- ♦ Yamashita J, Ogawa M, Egami H, Matsuo S, Kiyohara H, Inada K *et al.* Abundant expression of immunoreactive endothelin-1 in the growth of stromal cells in phyllodes tumour. *Cancer Res.* 1992; 52(14): 4046-9.
  - ♦ Yanagisawa M, Kurihara H, Kimura S, Tomobe Y, Kobayashi M, Mitsui Y *et al.* A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 1988; 332: 411-5.
  - ♦ Yanagisawa M, Masaki T. Molecular biology and biochemistry of the endothelins. *Trends Pharmacol Sci* 1989, 10, 374-9.
  - ♦ Yokokawa K, Mandal AK, Kohno M, Horio T, Murakawa K, Yasunari K, Takeda T. Heparin suppresses endothelin-1 action and production in spontaneously hypertensive rats. *Am J Physiol.* 1992 Nov; 263(5 Pt 2): R1035-41.
  - ♦ Zaho Y, Sprigall D, Hamid Q, Levene M, Polak J. Localisation and characterisation of endothelin-1 receptor binding in the blood vessels of human pulmonary tumours. *J. Cardiovasc. Pharmacy* 1995, 26(suppl.3), S341-5.
  - ♦ Zeiher AM, Ihling C, Pistorius K, Schachinger V, Schaefer H-E. Increased tissue endothelin immunoreactivity in atherosclerotic lesions associated with acute coronary syndrome. *Lancet* 1994, 344, 1405-6.
  - ♦ Zhao Y, Sprigall D, Hamid Q, Levene M, Polka J. Localisation and characterisation of endothelin-1 receptor binding in the blood vessels of human pulmonary tumours. *J. Cardiovasc. Pharmacy* 1995; 26(Suppl.3): S341-5.

- ♦ Ziche M, Morbidelli M, Donnini S, Ledda F. ETB receptors promote proliferation and migration of endothelial cells. *J Cardiovasc Pharmacol* 1995; 26(suppl 3): S284-6.
- ♦ Zonnenberg BA, Groenewegen G, Janus TJ, Leahy TW, Humerickhouse RA, Isaacson JD, Carr RA, Voest E. Phase I dose-escalation study of the safety and pharmacokinetics of atrasentan: an endothelin receptor antagonist for refractory prostate cancer. *Clin Cancer Res.* 2003 Aug 1; 9(8): 2965-72.
- ♦ Zuccarello M, Lewis AI, Rapoport RM. Endothelin ETA and ETB receptors in subarachnoid hemorrhage-induced cerebral vasospasm. *Eur J Pharmacol* 1994 Jun 23; 259(1): R1-2.